CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 38, 2009

A

Abbey, H. M., 35
Adams, Aaron S., 1022
Adams, A. S., 67
Adams, S. M., 1022
Addesso, K. M., 216
Adler, L. S., 207
Adler, P. H., 930
Afkhami, M. E., 1086
Aghdam, H. R., 885
Agnello, A. M., 433
Aguiar, C.m.l., 1613
Albajes, R., 944, 1533
Allen, M. L., 551
Allgood, D. W., 657
Almeida, R.P.P., 103
Alonso, D., 11
Alston, D. G., 955, 1126
Aluja, M., 588
Alyokhin, A. V., 846
Andersen, P. C., 183
Andow, D. A., 293, 1528
Anger, W., 870
Anulewicz, A. C., 1668
Arbogast, R. T., 561
Arpaia, S., 293, 1528
Ashfaq, M., 1276
Asís, J. D., 608
Atapour, M., 260
Aubert, M., 514
Aukema, B. H., 1189

B

Babcock, J., 281
Bachmann, A. C., 1347
Bagley, M. J., 1312
Bahlai, C. A., 708, 1106
Baker, T. C., 1745
Balentine, K. M., 1145
Barthell, J. F., 484
Baum, K. A., 1411
Beachy, J. A., 1226
Bean, D. W., 1373
Bell, H. A., 1533
Bennett, A., 93
Berberet, R. C., 1765
Bergh, J. C., 530, 1219
Bextine, B., 1360
Biddinger, D. J., 935
Biedermann, P.H.W., 1096
Bigelow, C. A., 996
Bigler, F., 1533
Bilu, E., 153
Bispo, P. C., 1613
Blackmer, J. L., 159
Blossey, B., 1653
Boina, D. R., 1250
Boiteau, G., 168
Bosque-Pérez, N. A., 836, 1429
Bossart, J. L., 43
Bouchard, R. W., Jr, 250
Bougeard, S., 514
Bowman, D. M.J.S., 1013
Boyd, B. M., 1211
Boykin, D., 551
Bradshaw, C.J.A., 1013
Brewster, C. C., 677, 1716
Brightwell, R. J., 1618
Brodbeck, B. V., 183
Brook, B. W., 1013
Brown, R. L., 815
Bruce, A. Y., 904
Brunke, A. J., 1106
Brust, M. L., 1521
Buchman, N., 962
Buddle, C. M., 797
Burgess, E.P.J., 1533
Burkle, L., 1293
Burkness, E. C., 1563
Burrack, H. J., 1400
Buteler, M., 1707
Byers, J. A., 159

C

Caldera, E. J., 78
Cameron, E. A., 1, 513
Campbell, J. F., 715
Cañas, L., 129
Cappaert, D., 1668
Carey, D. R., 1539
Carlson, J. E., 686
Carpentier, P., 514
Carver, S., 1585
Castello, J. D., 1235
Ceballos, R., 781
Celaya, R., 985
Center, T. D., 1145
Cerdá, X., 317
Chanbusarakum, L. J., 1069
Chandler, M. T., 823
Chapman, A. V., 677
Chauzat, M.-P., 514
Chen, C. C., 715
Chen, F. J., 26
Chen, L., 1480
Chen, Y., 1756
Cheng, L. L., 715
Chirino, M. G., 198
Chiroleu, F., 896
Chouinard, G., 870
Christian, C. E., 325
Clarke, S. R., 472
Coates, B. S., 1312
Cohen, M., 137
Colinet, H., 242
Coll, M., 153
Connell, J. H., 1400
Cooper, R. J., 1161
Cooper, W. R., 417
Cormier, D., 870
Corn, J. G., 971
Costamagna, A. C., 1343
Cottrell, T. E., 411
Cougoule, N., 514
Covelli, J., 505
Cronin, J. T., 1411
Crous, P. W., 143
Cruz-López, L., 1423
Cuda, J. P., 1642
Cuddington, K., 962
Currie, C. R., 78

D

Daane, K. M., 1485, 1496
D'Adamo, P., 803
Dadour, I. R., 633
Dalin, P., 1373
Daniels, J. C., 1211
Daugherty, M. P., 103, 667
Davis, R. S., 2
Davis, T. S., 639
de la Giroday, H.-M. C., 1189
Delfino, M. A., 505
de Little, S. C., 1013
Dillard, H. R., 1347
Dilling, C., 53
Ding, H., 1429
Ding, J., 1135, 1653
Dively, G. P., 1533
Dorhout, D. L., 274
Drapeau, P., 797
Dray, F. A., Jr, 1145
Dreyer, L. L., 143
Dudley, T., 1373

E

Edward, D. A., 1608
Egetter, D., 1161
Eigenbrode, S. D., 836, 1429
Elliott, S. E., 748, 1680
El Mujtar, V., 505
Epsky, N. D., 1259
Espinoza, F., 333
Esquivel, J. F., 766
Esquivel, S. V., 766
Etheridge, J. L., 35
Evans, A., 1241

Evans, E. W., 1283
Evans, T. A., 35
Evenden, M. L., 459

F
Faccoli, M., 307
Fathipour, Y., 885
Faucon, J.-P., 514
Faust, L. F., 1505
Feder, J. L., 588
Federici, B. A., 1533
Fernandes, O. A., 2
Ferrington, L. C., Jr, 250
Ferry, N., 1533
Finlayson, C. J., 846
Fleischer, S. J., 935, 1347, 1546
Folgarait, P. J., 198
Follett, P. A., 103, 1168
Forbes, A., 588
Forister, M. L., 1028
Fornell, A. M., 1400
Franc, A., 569
Frank, D. L., 1219
Frantz, J. M., 129
Fraser, I., 1226
Fraulo, A. B., 137
Frewin, A., 708
Friedrich, R., 600
Funderburk, J., 333, 1205

G
Galvan, T. L., 1563
Gao, J.-R., 499
Gao, Y., 1336
García, R. R., 985
García, U., 985
Garcia, C. M., 478
Gatehouse, A.M.R., 1533
Gayubo, S. F., 608
Ge, F., 26
Geden, C. J., 700
Gehring, C., 93
Geib, S. M., 686
Gilbert, L. E., 198
Gilburn, A. S., 1608
Gildow, F. E., 1347
Giles, K. L., 1765
Gillmore, J. L., 1642
Ginzel, M. D., 425
Godfrey, L. D., 1737
Golden, M., 103, 1168
Gómez, A. A., 11
González, J. A., 608
González-Hernández, H., 1423
Gotoh, T., 1626
Gould, F., 1336
Gourdon, F., 896
Grant, J., 53
Grau, O., 505
Gray, D. R., 18
Greco, N. M., 380
Greene, J. K., 450
Grenkow, L., 1467
Grey, W. E., 387
Gries, G., 1395
Griffith, J. L., 484
Groves, R. L., 1485
Gruber, M. Y., 1467
Gu, H., 524
Guedes, R.N.C., 1078
Guillén, L., 588
Gut, L. J., 955

H
Habeck, D. H., 1642
Hackenberger, B. K., 1600
Hagen, S. B., 627
Hall, D. G., 225
Hallett, R. H., 708, 1106, 1387
Hamm, R. L., 499
Han, Y., 1182
Hance, T., 242
Handelsman, J., 67
Hanif, M. A., 1276
Hanks, L. M., 395, 425, 651, 1267, 1462, 1595
Hansen, N. M., 627
Hanula, J. L., 1005
Harestad, A. S., 823
Harris, M. K., 1690
Hartman, G. L., 110
Hazzard, R. V., 207
Heath, R. N., 356
Heath, R. R., 1259
Hegedus, D. D., 1467
Hellmich, R. L., 1312, 1533
Henderson, G., 1241
Henne, D. C., 539
Herbert, D. A., Jr, 375, 1716
Herbert, J. J., 1155
Herron, G., 1575
Higgins, L. S., 281
Higley, L. G., 2
Hix, R. L., 1345
Hoback, W. W., 1521
Hoddle, M., 1634
Hofstetter, R. W., 639
Holben, W. E., 1022
Holtzer, T. O., 1061
Hooks, C.R.R., 442
Hoover, K., 686, 1745
Horton, D. R., 815
Hou, M., 1182
Hranitz, J. M., 484
Hsu, C. L., 433
Hua, K. H., 879
Huang, R. N., 879
Hunter, M. D., 1161
Hunter, W. B., 183
Hutchison, W. D., 1563

I
Imai, K., 404
Ims, R. A., 627
Irvin, N., 1634
Irwin, R., 1293
Isaacs, R., 955
Ito, M., 1324
Ivanov, K., 1724

J
Jackson, H. B., 1411
Jarić, D., 1600
Jia, F.-X., 1365
Jimenez-Gasco, M.D.M., 686
Jiménez-Martínez, E. S., 836
Jin, L., 1480
Johnson, M. W., 1485, 1496
Johnson, S. D., 741
Johnson, S. J., 539
Jones, B. C., 459
Joshi, A., 1539
Joyce, B. L., 1161
Jumean, Z., 1395
Jung, J. K., 1301
Júregui, B. M., 985

K
Kain, D. P., 530
Kajimura, H., 1324
Kajita, Y., 1283
Kalcounis-Rueppell, M. C., 657
Kamminga, K. L., 375, 1716
Kang, S.-T., 1301
Kard, B., 35
Kawai, S., 1626
Keena, M. A., 1745
Keiper, J., 1724
Kendra, P. E., 1259
Khan, M. I., 1276
Kim, K. S., 1312
Klein, K. T., 1555
Klepzig, K. D., 67, 1096
Klingeman, W. E., 530
Koch, R. L., 1563
Koopmans, J. M., 1189
Koppel, A. L., 375
Krčmar, S., 1600
Krupke, C. H., 176
Kuhar, T. P., 375, 677, 1716

L
Lachaize, J., 514
Lai, L. C., 879
Lambdin, P., 53
Landolt, P. J., 815
Latham, D. R., 1117
Lauzon, C. R., 116
Layton, R. J., 281
Leblanc, L., 1446
Lee, C.-Y., 1697
Lenz, M., 35
Leonard, D. S., 1555
Leskey, T. C., 168, 530, 870, 1219
Leslie, T. W., 935
Letkowski, S., 1235

Levac, J., 1387
Li, X., 1467
Li, Xiaoxia, 1480
Liburd, O. E., 137
Lin, G. G.-H., 499
Lin, W., 1182
Lindgren, B. S., 1189
Liu, D. L., 1575
Liu, H., 1480
Liu, T.-X., 1661
Lizama, M., 781
Locke, J. C., 129
Louda, S. M., 731
Lövei, G. L., 293, 1528
Lozada, M., 803
Lu, C. T., 715
Lumbierres, B., 944
Luna, J. M., 809
Luna, M. G., 365
Luo, T.-H., 1053
Luong-Skovmand, M. H., 569
Luttrell, R. G., 450

M

MacCollom, G. B., 116
Madec, F., 514
Malo, E., 861
Malone, L. A., 1533
Manandhar, R., 442
Mao, L., 1241
Marco, V., 1513
Marcum, D. B., 1737
Mareggiani, G. S., 380
Mares-Guia, T. R., 1078
Margolies, D. C., 715
Marois, J. J., 1205
Marquardt, P. T., 176
Marshall, J. M., 1226
Martel, A.-C., 514
Mastro, V. C., 1226
Matsumoto, Y., 1626
Matusick, G., 1235
Mauldin, J. K., 35
McAuslane, H. J., 216, 1155
McCullough, D. G., 1668
McKee, F. R., 1387
Meagher, R. L., 1546
Medal, J. C., 1642
Medina-Ortega, K. J., 836
Meloche, F., 168
Merrill, S. C., 1061
Meyer, Wendy L., 1250
Meyer, W. L., 116
Mian, M.A.R., 1301
Michel, A. P., 1301
Millar, J. G., 1267, 1462
Miller, D. A., 657
Miller, N. J., 274
Miller, T.E.X., 731
Mills, N. J., 1117
Mitchell, R. F., 395
Mizell, R. F., 1038
Mizell, R. F., III, 1155
Mizell, Russell F., III, 183
Mo, J., 1575
Moellenbeck, D. J., 281
Moharramipour, S., 260
Montgomery, W. S., 1259
Moon, R. D., 700
Moreira, J. A., 1462
Moreno, F., 1513
Morgan, D.J.W., 1485
Morrell, J. J., 478
Morrill, W. L., 387
Mound, L., 333
Mowry, T. M., 1429
Mueke, J., 904
Mullin, C. A., 935
Muñiz, M., 11
Muñoz-Ledesma, F. J., 723
Mutis, A., 192, 781
Myers, H. M., 930

N

Nadel, H., 1496
Nagoshi, R. N., 1546
Nam, Y., 121
Naranjo, S. E., 1533
Nault, B. A., 1347
Nechols, J. R., 715
Neese, P., 281
Nehme, M. E., 1745
Neoh, K.-B., 1697
Ngumbi, E., 836
Ni, X., 411
Niño, E. L., 493
Noda, H., 1626
Nombela, G., 11
Northfield, T. D., 1038
Northfield, Tobin D., 183

O

O'Brien, C. W., 731
O'Connell, N. V., 1400
Ohba, S.-Y., 856
Ohsaki, N., 404
Ohtsuka, K., 920
Olday, F., 116
Oliveira, M.G.A., 1078
Olson, D. L., 1539
Onagbola, E. O., 1250
O'Neal, M. J., 1373
Onyilagha, J., 1467
Opuni-Frimpong, E., 43
Osakabe, M., 920
Osekre, E. A., 1205
Osoro, K., 985
Overall, L. M., 484

P

Paini, D. R., 183
Palacios, R., 317
Palma, R., 192
Pardo, F., 192, 781
Parra, L., 192, 781
Payton, M. E., 1765
Peairs, F. B., 1061
Pedrosa-Macedo, J. H., 1642
Pence, J. A., 1211
Peng, G., 1439
Pereault, R. J., 1126
Pereyra, P. C., 365
Pérez-Moreno, I., 1513
Perfecto, I., 790
Perich, F., 192, 781
Peterson, R.K.D., 2, 1707
Pfannenstiel, R. S., 1379
Phillips, P. A., 1400
Philpott, S. M., 600, 790
Pichette, A., 870
Poland, T. M., 757, 1668, 1756
Pons, X., 944
Porta, P., 514
Porter, E. W., 846
Poulsen, M., 78
Powell, B. E., 1618
Prado, S. S., 103
Pratt, P. D., 1145
Purcell, M. F., 1135
Pureswaran, D. S., 757

Q

Quesada-Moraga, E., 723
Quilici, S., 896
Quiroz, A., 192, 781

R

Radjabi, G., 885
Raffa, K. F., 67
Ranger, C. M., 129
Ray, A. M., 425, 1462
Rayamajhi, M. B., 1145
Reding, M. E., 129
Reisig, D. D., 1737
Reissig, W. H., 433
Retana, J., 317
Rezapanah, M., 885
Rhea, R., 53
Ribon, A.O.B., 1078
Rice, M. E., 274
Richardson, M. L., 651, 1595
Richmond, D. S., 996
Riddle, T. C., 183, 1038
Rieske, L. K., 417
Riggio, M. R., 1174
Ripa, R., 333
Robert, T., 1411
Rocca, M., 380
Rodriguez, F., 333
Rodriguez-Saona, C., 861
Roets, F., 143
Roitberg, B. D., 823
Rojas, J. C., 1423
Roland, J., 459
Romeis, J., 1533
Romero, C. M., 1267

Roubaud, M., 896
Roush, R., 1533
Rousse, P., 896
Roux, J., 356
Rubinoff, D., 1446
Rudgers, J. A., 1086
Ruiz-Montiel, C., 1423
Rull, J., 588
Rutledge, C. E., 1267
Ryoo, M. I., 121

S
Sackett, W. M., 1347
Saint-Germain, M., 797
Sánchez, N. E., 365
Sánchez-Martínez, G., 1197
Santiago-Álvarez, C., 723
Santos, G. M. de M., 1613
Sappington, T. W., 274, 1312
Schaafsma, A. W., 708
Schulthess, F., 904
Schultz, P. B., 677
Scott, J. G., 499
Sears, M. K., 708, 1106, 1533
Sehnal, F., 1533
Senger, S. E., 823
Shah, D. A., 1347
Shelton, A. M., 1533
Shields, K. S., 577
Shuttleworth, A., 741
Silverman, J., 1618
Singh, A. P., 129
Sisterson, M. S., 345
Sivinski, J., 588
Sjogren, R. E., 116
Smith, J. F., 450
Smith, W. A., 551
Son, Y., 1485
Song, Y., 1480
Sorenson, C. E., 493
Soroka, J. J., 1467
Spafford, H., 1585
Stafford, K. C., III, 977
Stehman, S. V., 1235
Stelinski, L. L., 225, 861, 1250
Stevens, M., 1575
Stevenson, D. E., 1690
Stewart, K. M., 1608
Stodola, K. W., 1161
Storer, A. J., 1226
Storer, N., 281
Storey, A., 1585
Story, J. M., 971
Stout, M. J., 1174
Straight, K. E., 1347
Suarez-Espinoza, J., 1634
Suen, G., 78
Sun, J., 472
Sun, Y. C., 26
Sun, Z., 387
Swift, I. P., 1462
Szendrei, Z., 861

T
Taborsky, M., 1096
Takahashi, M., 731
Tanaka, K., 235, 266
Taylor, R. M., 1379
Teal, P.E.A., 561
Teale, S. A., 1235
Teixeira, L.A.F., 955
Thistlewood, H.M.A., 823
Thorp, R. W., 484
Tien, M., 686
Tillman, P. G., 1038
Tingle, C., 450
Tobin, P. C., 1555
Tomberlin, J. K., 930
Tormos, J., 608
Torto, B., 561
Trager, M. D., 1211
Trotter, R. T., III, 577
Tufts, D. M., 1360
Tyson, R., 823

U
Ullman, D. E., 1069
Ulyshen, M. D., 1005
Underwood, E. C., 325
Uno, S., 790

V
Vandermeer, J., 790
Van Wyk, M., 356
Vargas, R. I., 1446
Vincent, C., 168
Visôtto, L. E., 1078
Vogt, J. T., 551
Vorsa, N., 129
Voss, S. C., 633
Vossen, P. M., 1400

W
Wagner, M. R., 1197
Walgenbach, J. F., 530
Wallet, B., 551
Wang, J.-J., 1365
Wang, X.-G., 1496
Wang, Xiaojuan, 1480
Wang, Xu, 1439
Wang, Y., 1135
Ward, J. S., 977
Washburn, S. P., 493
Watson, D. W., 493
Weaver, D. K., 387, 1707
Weinstein, P., 1585
Wenda-Piesik, A., 387
Wenninger, E. J., 225
Werner, B. J., 1429
Westcott, N. D., 1467
Weston, P. A., 1505
Whalon, M. E., 1126
Wharton, R., 588
Wheeler, G. S., 1135
Whelan, P. I., 1013
White, L. J., 971
Wiedenmann, R. N., 1344
Wille, B. D., 110
Williams, S. C., 977
Willms, S., 561
Wingfield, M. J., 143, 356
Wolt, J. D., 1533
Wood, B. W., 411
Wood, C., 1395
Worthley, T. E., 977
Wright, D. L., 1205
Wright, M. G., 442, 1168
Wright, R. J., 1521
Wright, S. E., 870
Wu, K., 1336
Wu, W. J., 879

X
Xie, Y., 1439
Xu, N., 1467
Xue, J., 1439
Xue, L., 809
Xue, Y., 708

Y
Yamanaka, T., 235
Yang, C. C., 879
Yang, M.-S., 1365
Yang, P. S., 715
Yang, W.-J., 1365
Yang, X., 1241
Yang, X.-B., 1661
Yang, Y., 1174
Yasuda, H., 1283
Yoon, E. Y., 121
Yu, X.-D., 1053

Z
Zack, R. S., 815
Zalom, F. G., 1400
Zarrabi, A. A., 1765
Zehnder, C. B., 1161
Zhang, A., 530, 1745
Zhang, Aijun, 870
Zhang, G., 1135
Zhang, L., 472
Zhang, W., 1301
Zhang, Y., 1439
Zhou, H.-Z., 1053

CUMULATIVE KEY WORD INDEX
Environmental Entomology
Volume 38, 2009

A

abdomen length,
alkaloids, 879
body length, 879
Solenopsis geminata, 879
venom volume, 879
Acalymma vittatum,
Erwinia tracheiphila, 395
frass, 395
polymerase chain reaction, 395
vector–pathogen interactions, 395
Acrobasis nuxvorella,
circadian, 1690
pecan nut casebearer, 1690
periodicity, 1690
pheromone, 1690
Acrosternum hilare, 450, 1716
black light trap, 1716
degree day, 1716
Euschistus servus, 450
flight activity, 1716
Glycine max, 450
Nezara viridula, 450
population dynamics, 1716
trap crop, 450
Acyrthosiphon pisum,
Aphis glycines, 1347
Aphis gossypii, 1347
Phaseolus vulgaris, 1347
Therioaphis trifolii, 1347
Adelges tsugae,
climate, 577
hemlock, 577
invasive species, 577
landscape, 577
adhesives,
behavior, 168
bonding, 168
cyanoacrylate, 168
ethological software, 168
Aedes vigilax,
Culex annulirostris, 1013
density dependence, 1013
larval habitats, 1013
vector control, 1013
Aegorhinus superciliosus,
contact pheromone, 192
mating behavior, 192
raspberry weevil, 192
Aeromonas salmonicida,
disease-pathogen dynamics, 1022
insect feeding assay, 1022
Plecoptera, 1022
trophic interactions, 1022
Agapeta zoegana,
biological control, 971
Centaurea stoebe, 971
Cyphocleonus achates, 971
diapause, 971
larval development, 971
agave,
2-methyl-4-octanone, 1423
pheromone release, 1423
weevil, 1423
age,
dispersal potential, 235
flight mill, 235
photoperiod, 235
ragweed beetle, 235
age-dependent developmental rate,
egg phenology, 18
phenology, 18
temperature, 18
aggression,
interspecific competition, 425
longhorned beetle, 425
mating behavior, 425
Agral 90,
antixenosis, 1387
pea leafminer, 1387
preoviposition behaviors, 1387
Sylgard 309, 1387
agricultural landscapes,
biological control, 380
blueberry, 380
cottony cushion scale, 380
risk of parasitism, 380
Agrilus planipennis,
emerald ash borer, 1226
Fraxinus, 1226
trap tree, 1226
tree stress, 1226
agroecology,
agroecosystem, 1028
cross-boundary, 1028
Great Basin, 1028
spillover, 1028
agroecosystem,
agroecology, 1028
cross-boundary, 1028
Great Basin, 1028
spillover, 1028
alkaloids,
abdomen length, 879
body length, 879
Solenopsis geminata, 879
venom volume, 879
allocation,
foraging, 1379
Helicoverpa zea, 1379
nutrition, 1379
starvation, 1379
ambrosia beetle,
cytochrome oxidase I, 1324
genetic structure, 1324
nested clade phylogeographic analysis, 1324
ammonia,
biosolid cake, 633
fly production, 633
temperature, 633
Anaphothrips obscurus,
economic threshold, 1737
hybrid economic injury level, 1737
multilinear regression, 1737
angiosperm volatiles,
Engelmann spruce weevil, 1189
nonhost volatiles, 1189
plant–insect interactions, 1189
spatial point processes, 1189
Anisopteromalus calandrae,
Aspergillus candidus, 121
Aspergillus niger, 121
positive interaction, 121
Sitophilus oryzae, 121
Anoplophora glabripennis, 686, 1745
Cerambycidae, 1745
culture-independent community analysis, 686
gut microbiota, 686
host tree resistance, 686
male-produced pheromone, 1745
mate-finding, 1745
trap design, 1745
ant,
coccinellid, 846
native, 846
natural enemies, 846
non-native, 846
ant community,
citrus orchard, 317
dominance, 317
Mediterranean region, 317
seasonality, 317
ant nest,
nest-site limitation, 600
urban ecology, 600

antagonism,
Apocynaceae, 741
mutualism, 741
nectar, 741
pollination syndromes, 741
ant–aphid interaction,
ant–aphid interaction, 1618
foraging, 1618
precipitation, 1618
Anthonomus eugenii,
Capsicum annuum, 216
host plant volatiles, 216
Solanum americanum, 216
Solanum melongena, 216
antixenosis,
Agral 90, 1387
pea leafminer, 1387
preoviposition behaviors, 1387
Sylgard 309, 1387
Anura,
diving beetle, 856
fish, 856
Odonata, 856
predator, 856
aphid behavior,
emigration, 1429
Myzus persicae, 1429
potato, 1429
virus-induced volatiles, 1429
aphid population fitness,
green peach aphid, 129
Myzus persicae, 129
peroxidase, 129
phenolics, 129
Aphidius colemani,
aphids, 153
Coccinella undecimpunctata, 153
Myzus persicae, 153
prey quality, 153
aphid–plant interaction,
aphid–plant interaction, 411
Carya illinoinensis, 411
host finding, 411
leaf chlorosis, 411
Melanocallis caryaefoliae, 411
aphids,
Aphidius colemani, 153
Coccinella undecimpunctata, 153
Myzus persicae, 153
prey quality, 153
Aphis craccivora,
cowpea aphid, 1765
developmental threshold, 1765
within-plant patterns, 1765
Aphis glycines, 1301, 1347
Acyrthosiphon pisum, 1347
Aphis gossypii, 1347
microsatellites, 1301
Phaseolus vulgaris, 1347
population genetic structure, 1301
soybean aphid, 1301
Therioaphis trifolii, 1347
Aphis gossypii,
Acyrthosiphon pisum, 1347
Aphis glycines, 1347
Phaseolus vulgaris, 1347
Therioaphis trifolii, 1347
Aphthona flea beetles,
chill degree-days, 1539
low temperature, 1539
overwintering, 1539
snow cover, 1539
Apis mellifera, 207, 514
Bombus spp., 207
colony health, 514
field survey, 514
France, 514
multispecies interactions, 207
Peponapis pruinosa, 207
pesticides, 514
winter squash, 207
Apocynaceae,
antagonism, 741
mutualism, 741
nectar, 741
pollination syndromes, 741
apple, 530, 1219
behavior, 1219
dogwood borer, 1219
forest, 530
ornamentals, 530
pheromone trapping, 530
Sesiidae, 1219
Synanthedon scitula, 530, 1219
arachnids,
breeds, 985
grazing pressure, 985
insects, 985
ungulates, 985
Araneidae,
Argiope trifasciata, 651
Cyclosa turbinata, 651
Mangora gibberosa, 651
Neoscona arabesca, 651
arbovirus,
community, 1585
dryland salinity, 1585
secondary salinization, 1585
temporary water, 1585
arbuscular mycorrhiza,
ectomycorrhiza, 93
insect herbivory, 93
insect pollination, 93
Argentina,
biological control, 198
fire ants, 198
parasitoid, 198
social form, 198
Argiope trifasciata, 651, 1595
Araneidae, 651
Cyclosa turbinata, 651, 1595
Mangora gibberosa, 651, 1595
Neoscona arabesca, 651, 1595
species diversity, 1595
Arribes del Duero Natural Park,
biogeography, 608
diversity, 608
Mediterranean region, 608
solitary wasps, 608
Arsenophonus,
bacterial symbiont, 110
Buchnera aphidicola, 110
soybean aphid, 110
artificial medium,
biological invasion, 1096
cooperative breeding, 1096
insect agriculture, 1096
symbiosis, 1096
Asian citrus psyllid,
attractant, 225
citrus greening disease, 225
kairomones, 225
olfaction, 225
Aspergillus candidus,
Anisopteromalus calandrae, 121
Aspergillus niger, 121
positive interaction, 121
Sitophilus oryzae, 121
Aspergillus niger,
Anisopteromalus calandrae, 121
Aspergillus candidus, 121
positive interaction, 121
Sitophilus oryzae, 121
Attini,
Escovopsis, 78
Leucoagaricus, 78
Pseudonocardia, 78
symbiosis, 78
attractant, 225, 870
Asian citrus psyllid, 225
citrus greening disease, 225
electroantennogram, 870
gas chromatographic-electroantennographic detection, 870
kairomones, 225
olfaction, 225
pheromone, 870
weevil, 870
Aucuba japonica,
gall insect, 404
physical defense, 404
plant–herbivore interaction, 404
susceptible period, 404
Austrocedrus chilensis,
Cinara cupressi, 505
Cinara tujafilina, 505
cytochrome oxidase I, 505
polymerase chain reaction-restriction fragment length polymorphism, 505
automated trapping,
flight behavior, 1555
Lymantria dispar, 1555

phenology, 1555
sampling, 1555
autumnal moth,
pupal predation, 627
sub-arctic birch forest, 627
winter moth, 627

B

bacteria symbionts,
pest control, 1078
protease inhibitor, 1078
proteases, 1078
16S rRNA, 1078
bacterial interaction,
biological control, 1360
insect symbiont, 1360
bacterial symbiont, 110, 1069
Arsenophonus, 110
Buchnera aphidicola, 110
distribution, 1069
Frankliniella occidentalis, 1069
prevalence, 1069
soybean aphid, 110
Bactrocera,
Cue-Lure, 1446
methyl eugenol, 1446
Bactrocera oleae,
egg maturation, 1496
high temperature, 1496
reproduction, 1496
survival, 1496
baiting,
castes, 35
larvae, 35
reproductive biology, 35
subterranean termites, 35
bark beetle phermone,
cis-verbenol, 472
frontalin, 472
red turpentine beetle, 472
trans-verbenol, 472
bark beetles, 67, 1197
context dependency, 67
Dendroctonus rhizophagus, 1197
host preference, 1197
insects, 67
microorganisms, 67
pine seedlings, 1197
symbiosis, 67
tree mortality, 1197
Barley yellow dwarf virus epidemiology,
bird cherry-oat aphid, 836
plant–insect–virus interactions, 836
transgenic virus resistance, 836
virus-induced volatiles, 836
Beauveria bassiana,
Heterorhabdtis, 1126
inundative biological control, 1126
Metarhizium anisopliae, 1126
Steinernema, 1126
beech bark disease,
beech scale, 1235
Cryptococcus fagisuga, 1235
Fagus grandifolia, 1235
beech scale,
beech bark disease, 1235
Cryptococcus fagisuga, 1235
Fagus grandifolia, 1235
beetle,
bidirectional selection, 266
exotic insect, 266
realized heritability, 266
repeatability, 266
behavior, 168, 823, 1219
adhesives, 168
apple, 1219
bonding, 168
cyanoacrylate, 168
diffusion model, 823
dispersal, 823
dogwood borer, 1219
ethological software, 168
habitat selection, 823
perceptual range, 823
Sesiidae, 1219
Synanthedon scitula, 1219
Bemisia argentifolii,
intercropping, 442
living mulch, 442
phytotoxemia, 442
squash, 442
bess beetle,
coarse woody debris, 1411
horned passalus, 1411
landscape matrix, 1411
bidirectional selection,
beetle, 266
exotic insect, 266
realized heritability, 266
repeatability, 266
biodiversity,
coarse woody debris, 1005
colonization, 1005
dispersal, 1005
epigaeic, 1005
biogeography,
Arribes del Duero Natural Park, 608
diversity, 608
Mediterranean region, 608
solitary wasps, 608
biological control, 2, 198, 365, 380, 539, 677, 708, 715, 723, 896, 971, 1360, 1634
Agapeta zoegana, 971
agricultural landscapes, 380
Argentina, 198
bacterial interaction, 1360
blueberry, 380
Centaurea stoebe, 971
circadian rhythm, 896
climatic parameters, 896
cottony cushion scale, 380
Cyphocleonus achates, 971
diapause, 971
dispersal, 677
endophyte, 723
European corn borer, 677
feeding behavior, 715
fire ants, 198
functional response, 708, 1634
Gonatocerus ashmeadi, 1634
Gonatocerus tuberculifemur, 1634
Homalodisca vitripennis, 1634
host selection, 896
host–parasitoid interaction, 365
insect demography, 2
insect symbiont, 1360
integrated pest management, 677
larval development, 971
Lycopersicon esculentum, 365
multicolored Asian lady beetle, 708
multiple decrement life table, 2
natural enemies, 2
parasitoid, 198
phenology, 539
physiological state, 896
phytophagous mites, 715
poppy stem gall wasp, 723
predator–prey interaction, 715
predatory potential, 715
relative abundance, 539
risk analysis, 2
risk of parasitism, 380
sampling, 539
seven-spotted lady beetle, 708
social form, 198
Solenopsis invicta, 539
soybean aphid, 708
spatial scales, 365
systemic protection, 723
Timaspis papaveris, 723
Trichogramma, 677
biological control of weeds,
Cirsium, 731
ecological risk assessment, 731
Trichosirocalus horridus, 731
weed control, 731
biological invasion, 274, 484, 1096
artificial medium, 1096
cooperative breeding, 1096
insect agriculture, 1096
NADH dehydrogenase 1, 274
nest site selection, 484
parasitoid, 484
stress, 484
Striacosta albicosta, 274
symbiosis, 1096
temperature, 484

Zea mays, 274
biological weed control,
China, 1135
host specificity, 1135
invasive plant, 1135
leaf-rolling weevil, 1135
biosafety,
laboratory experiments, 293
natural enemies, 293
review, 293
transgenic plants, 293
biosolid cake,
ammonia, 633
fly production, 633
temperature, 633
bird cherry-oat aphid,
Barley yellow dwarf virus epidemiology, 836
plant–insect–virus interactions, 836
transgenic virus resistance, 836
virus-induced volatiles, 836
black light trap,
Acrosternum hilare, 1716
degree days, 1716
flight activity, 1716
population dynamics, 1716
black soldier fly,
development, 930
Hermetia illucens, 930
temperature, 930
blacklegged tick,
Japanese barberry, 977
Lyme disease, 977
white-footed mouse, 977
white-tailed deer, 977
blueberry,
agricultural landscapes, 380
biological control, 380
cottony cushion scale, 380
risk of parasitism, 380
body length,
abdomen length, 879
alkaloids, 879
Solenopsis geminata, 879
venom volume, 879
Bombus appositus,
bumble bee, 1680
Delphinium barbeyi, 1680
flower density, 1680
food limitation, 1680
Bombus spp.,
Apis mellifera, 207
multispecies interactions, 207
Peponapis pruinosa, 207
winter squash, 207
bonding,
adhesives, 168
behavior, 168
cyanoacrylate, 168
ethological software, 168
Braconidae,
co-evolution, 588
North America, 588
Rhagoletis pomonella, 588
Tephritidae, 588
Brassica oleracea L. variety *italica* Plenck,
development, 11
nymphal survival, 11
phenological models, 11
Brazilian peppertree,
Christmas berry, 1642
natural enemies, 1642
risk assessment, 1642
breeds,
arachnids, 985
grazing pressure, 985
insects, 985
ungulates, 985
brown stink bug,
distribution, 1038
raster map, 1038
southern green stink bug, 1038
Bt,
Carabidae, 935
Cry3Bb, 935
Harpalus pensylvanicus, 935
neonicotinoids, 935
Bt corn, 176, 281
Cry1F, 281
dispersal, 176
field monitoring, 281
insect resistance management, 176
nontarget arthropods, 281
western corn rootworm, 176
Bt cotton,
Cry1Ac, 1336
Helicoverpa armigera, 1336
resistance frequency, 1336
Buchnera,
Heteroptera, 103
southern green stink bug, 103
symbiosis, 103
Buchnera aphidicola,
Arsenophonus, 110
bacterial symbiont, 110
soybean aphid, 110
bumble bee, 748, 1680
Bombus appositus, 1680
Delphinium barbeyi, 1680
flower density, 748, 1680
food availability, 748
food limitation, 1680
foraging distance, 748
subalpine, 748
Buprestidae,
Coleoptera, 1756
girdling, 1756
insect–plant interactions, 1756
light intensity, 1756

C

C. transversoguttata,
Coccinella septempunctata, 1283
egg size, 1283
egg viability, 1283
reproduction, 1283
Caatinga,
community ecology, 1613
Polistinae, 1613
species richness, 1613
Cacopsylla pyricola,
diel periodicity, 815
pear psylla, 815
sex attraction, 815
canonical correlation analysis,
canopy trap, 1600
distributional pattern, 1600
elevational distribution, 1600
transect, 1600
canopy arthropods,
eastern hemlock, 53
hemlock woolly adelgid, 53
horticultural oil, 53
imidacloprid, 53
canopy trap,
canonical correlation analysis, 1600
distributional pattern, 1600
elevational distribution, 1600
transect, 1600
Capsicum annuum,
Anthonomus eugenii, 216
host plant volatiles, 216
Solanum americanum, 216
Solanum melongena, 216
carabid beetles,
corn, 944
genetically modified organisms, 944
Orius, 944
spiders, 944
Carabidae, 935, 1053
Bt, 935
Cry3Bb, 935
diversity, 1053
ecotones between regenerating and mature stands, 1053
edge-effects, 1053
Harpalus pensylvanicus, 935
neonicotinoids, 935
Caribbean fruit fly,
1,4-diaminobutane, 1259
olfaction, 1259
semiochemicals, 1259
synthetic attractants, 1259
Carya illinoinensis,
aphid–plant interaction, 411
host finding, 411
leaf chlorosis, 411
Melanocallis caryaefoliae, 411

castes,
baiting, 35
larvae, 35
reproductive biology, 35
subterranean termites, 35
Centaurea stoebe,
Agapeta zoegana, 971
biological control, 971
Cyphocleonus achates, 971
diapause, 971
larval development, 971
Cephus cinctus,
insect pathogens, 387
plant pathogens, 387
Triticum aestivum, 387
wheat, 387
Cerambycidae, 797, 1745
Anoplophora glabripennis, 1745
host selection, 797
male-produced pheromone, 1745
mate-finding, 1745
primary attraction, 797
Scolytinae, 797
semiochemicals, 797
trap design, 1745
Ceroplastes japonicus Green,
Chilocorus kuwanae Silvestri, 1439
methyl jasmonate, 1439
α-pinene, 1439
volatiles, 1439
chill degree-days,
Aphthona flea beetles, 1539
low temperature, 1539
overwintering, 1539
snow cover, 1539
Chilo suppressalis, 260, 1182
cold hardiness, 260
diapause, 260
glycerol, 260
overwintering, 1182
paddy rice, 1182
rice stem borer, 260
supercooling capacity, 1182
water-oat, 1182
Chilocorus kuwanae Silvestri,
Ceroplastes japonicus Green, 1439
methyl jasmonate, 1439
α-pinene, 1439
volatiles, 1439
China,
biological weed control, 1135
host specificity, 1135
invasive plant, 1135
leaf-rolling weevil, 1135
Choristoneura conflictana,
Malacosoma disstria, 459
monitor, 459
prediction, 459
sex pheromone traps, 459
Christmas berry,
Brazilian peppertree, 1642
natural enemies, 1642
risk assessment, 1642
chronological reproductive isolation,
flight phenology, 1697
Macrotermes, 1697
swarming, 1697
Chrysopa nigricornis,
field cage, 1117
field observation, 1117
Harmonia axyridis, 1117
laboratory arena, 1117
Cicadellidae,
feeding, 1485
glassy-winged sharpshooter, 1485
survival, 1485
temperature, 1485
vector, 1485
Cinara cupressi,
Austrocedrus chilensis, 505
Cinara tujafilina, 505
cytochrome oxidase I, 505
polymerase chain reaction-restriction fragment length polymorphism, 505
Cinara tujafilina,
Austrocedrus chilensis, 505
Cinara cupressi, 505
cytochrome oxidase I, 505
polymerase chain reaction-restriction fragment length polymorphism, 505
circadian,
Acrobasis nuxvorella, 1690
pecan nut casebearer, 1690
periodicity, 1690
pheromone, 1690
circadian rhythm,
biological control, 896
climatic parameters, 896
host selection, 896
physiological state, 896
Cirsium,
biological control of weeds, 731
ecological risk assessment, 731
Trichosirocalus horridus, 731
weed control, 731
cis-verbenol,
bark beetle phermone, 472
frontalin, 472
red turpentine beetle, 472
trans-verbenol, 472
citrus greening disease,
Asian citrus psyllid, 225
attractant, 225
kairomones, 225
olfaction, 225
citrus orchard,
ant community, 317
dominance, 317
Mediterranean region, 317
seasonality, 317
climate,
Adelges tsugae, 577
hemlock, 577
invasive species, 577
landscape, 577
climate change,
degree-day, 1505
firefly, 1505
phenology, 1505
synchronous display, 1505
climatic parameters,
biological control, 896
circadian rhythm, 896
host selection, 896
physiological state, 896
co-evolution,
Braconidae, 588
North America, 588
Rhagoletis pomonella, 588
Tephritidae, 588
coarse woody debris, 1005, 1411
bess beetle, 1411
biodiversity, 1005
colonization, 1005
dispersal, 1005
epigaeic, 1005
horned passalus, 1411
landscape matrix, 1411
Coccinella septempunctata,
C. transversoguttata, 1283
egg size, 1283
egg viability, 1283
reproduction, 1283
Coccinella undecimpunctata,
Aphidius colemani, 153
aphids, 153
Myzus persicae, 153
prey quality, 153
coccinellid,
ant, 846
native, 846
natural enemies, 846
non-native, 846
codling moth, 885, 1395
Cydia pomonella, 885, 1395
development, 885
frequency distribution, 1395
larval aggregation, 1395
linear and nonlinear models, 885
pheromone, 1395
thermal threshold, 885
coelopidae,
sex-biased parasitism, 1608
cold hardiness,
Chilo suppressalis, 260
diapause, 260
glycerol, 260

rice stem borer, 260
cold tolerance,
overwintering, 524
pest management, 524
Plutella xylostella, 524
Coleoptera,
Buprestidae, 1756
girdling, 1756
insect–plant interactions, 1756
light intensity, 1756
colonization,
biodiversity, 1005
coarse woody debris, 1005
dispersal, 1005
epigaeic, 1005
colonization process,
onion, 1575
population model, 1575
Thrips tabaci, 1575
colony health,
Apis mellifera, 514
field survey, 514
France, 514
pesticides, 514
community,
arbovirus, 1585
dryland salinity, 1585
secondary salinization, 1585
temporary water, 1585
community ecology,
Caatinga, 1613
Polistinae, 1613
species richness, 1613
community interactions,
endophyte, 417
gypsy moth, 417
lesser chestnut weevil, 417
Torymus, 417
competition, 26, 639
Dendroctonus brevicomis, 639
Dendroctonus frontalis, 639
elevated CO_2, 26
ponderosa pine, 639
population dynamics, 639
Rhopalosiphum padi, 26
Schizaphis graminum, 26
Sitobion avenae, 26
conservation management,
ecological disturbance, 325
mediterranean ecosystems, 325
restoration, 325
contact pheromone,
Aegorhinus superciliosus, 192
mating behavior, 192
raspberry weevil, 192
contaminant pest,
Harmonia axyridis, 1563
modeling, 1563
sugar content, 1563
trap catching, 1563
context dependency,
bark beetles, 67
insects, 67
microorganisms, 67
symbiosis, 67
context-dependent interactions,
Epichloë, 1086
food choice, 1086
Neotyphodium, 1086
Spodoptera frugiperda, 1086
cooperative breeding,
artificial medium, 1096
biological invasion, 1096
insect agriculture, 1096
symbiosis, 1096
Coptotermes formosanus,
foraging, 1241
integrated pest management, 1241
nesting, 1241
Solenopsis invicta, 1241
corn,
carabid beetles, 944
genetically modified organisms, 944
Orius, 944
spiders, 944
cotton fleahopper,
cotton pest, 766
host plant, 766
cotton pest,
cotton fleahopper, 766
host plant, 766
cottony cushion scale,
agricultural landscapes, 380
biological control, 380
blueberry, 380
risk of parasitism, 380
cowpea aphid,
Aphis craccivora, 1765
developmental threshold, 1765
within-plant patterns, 1765
crapemyrtle,
crapemyrtle aphid, 1155
host suitability, 1155
crapemyrtle aphid,
crapemyrtle, 1155
host suitability, 1155
cross-boundary,
agroecology, 1028
agroecosystem, 1028
Great Basin, 1028
spillover, 1028
crucifer flea beetle,
olfactometer bioassay, 1467
volatile compounds, 1467
Cry1Ac,
Bt cotton, 1336
Helicoverpa armigera, 1336
resistance frequency, 1336
Cry1F,
Bt corn, 281
field monitoring, 281
nontarget arthropods, 281
Cry3Bb,
Bt, 935
Carabidae, 935
Harpalus pensylvanicus, 935
neonicotinoids, 935
Cryptococcus fagisuga,
beech bark disease, 1235
beech scale, 1235
Fagus grandifolia, 1235
Cue-Lure,
Bactrocera, 1446
methyl eugenol, 1446
Culex annulirostris,
Aedes vigilax, 1013
density dependence, 1013
larval habitats, 1013
vector control, 1013
culture-independent community analysis,
Anoplophora glabripennis, 686
gut microbiota, 686
host tree resistance, 686
cuticular hydrocarbons,
5,17-dimethylnonacosane, 1267
9-heptacosyne, 1267
mating behavior, 1267
9-pentacosyne, 1267
cyanoacrylate,
adhesives, 168
behavior, 168
bonding, 168
ethological software, 168
Cyclosa turbinata, 651, 1595
Araneidae, 651
Argiope trifasciata, 651, 1595
Mangora gibberosa, 651, 1595
Neoscona arabesca, 651, 1595
species diversity, 1595
Cydia pomonella, 885, 1395
codling moth, 885, 1395
development, 885
frequency distribution, 1395
larval aggregation, 1395
linear and nonlinear models, 885
pheromone, 1395
thermal threshold, 885
Cylindroiulus caeruleocinctus,
integrated pest management, 1106
Pterostichus melanarius, 1106
Cyphocleonus achates,
Agapeta zoegana, 971
biological control, 971
Centaurea stoebe, 971
diapause, 971
larval development, 971
cytochrome oxidase,
haplotype, 1546
Noctuidae, 1546
Spodoptera frugiperda, 1546

cytochrome oxidase I, 505, 1324
ambrosia beetle, 1324
Austrocedrus chilensis, 505
Cinara cupressi, 505
Cinara tujafilina, 505
genetic structure, 1324
nested clade phylogeographic analysis, 1324
polymerase chain reaction-restriction fragment length polymorphism, 505
cytoplasmic incompatibility,
Laodelphax striatellus, 1626
microinjection, 1626
Nilaparvata lugens, 1626
Wolbachia, 1626

D

damage,
phenology, 307
precipitation, 307
temperature, 307
voltinism, 307
degree-day, 1505, 1521, 1716
Acrosternum hilare, 1716
black light trap, 1716
climate change, 1505
firefly, 1505
flight activity, 1716
grasshopper development, 1521
phenology, 1505, 1521
population dynamics, 1716
rangeland, 1521
synchronous display, 1505
Delphinium barbeyi,
Bombus appositus, 1680
bumble bee, 1680
flower density, 1680
food limitation, 1680
Dendroctonus brevicomis,
competition, 639
Dendroctonus frontalis, 639
ponderosa pine, 639
population dynamics, 639
Dendroctonus frontalis,
competition, 639
Dendroctonus brevicomis, 639
ponderosa pine, 639
population dynamics, 639
Dendroctonus rhizophagus,
bark beetles, 1197
host preference, 1197
pine seedlings, 1197
tree mortality, 1197
density dependence, 790, 962, 1013
Aedes vigilax, 1013
Culex annulirostris, 1013
environmental conditions, 962
fertilization, 962
host-parasitoid, 790
Identificaticoffee agroecosystem, 790
larval habitats, 1013
olfactory cue, 790
plant morphology, 962
reproduction, 962
scale-dependent effects, 790
vector control, 1013
detection methods,
girdled trees, 1668
Manuka oil, 1668
methyl jasmonate, 1668
trap tree, 1668
development, 11, 885, 930, 1061, 1661
black soldier fly, 930
Brassica oleracea L. variety *italica* Plenck, 11
codling moth, 885
Cydia pomonella, 885
fecundity, 1661
Hermetia illucens, 930
intrinsic rate of increase, 1061
linear and nonlinear models, 885
meta-analysis, 1061
nymphal survival, 11
oviposition, 1661
phenological models, 11
reproduction, 1061
Russian wheat aphid, 1061
Solanaceae, 1661
survival, 1661
temperature, 930
thermal threshold, 885
developmental period,
life cycle, 478
powderpost beetle, 478
thermal constants, 478
thermal requirements, 478
developmental threshold,
Aphis craccivora, 1765
cowpea aphid, 1765
within-plant patterns, 1765
Diabrotica undecimpunctata undecimpuncta,
integrated pest management, 809
sampling plan design, 809
scouting programs, 809
strip spraying, 809
1,4-diaminobutane,
Caribbean fruit fly, 1259
olfaction, 1259
semiochemicals, 1259
synthetic attractants, 1259
diapause, 260, 569, 971
Agapeta zoegana, 971
biological control, 971
Centaurea stoebe, 971
Chilo suppressalis, 260
cold hardiness, 260
Cyphocleonus achates, 971
glycerol, 260
larval development, 971
locust, 569
Madagascar, 569
reproductive maturity, 569
rice stem borer, 260
wing color, 569
Diaphorina citri,
dispersal behavior, 1250
huanglongbing, 1250
immunomarking, 1250
mark-capture, 1250
diel periodicity,
Cacopsylla pyricola, 815
pear psylla, 815
sex attraction, 815
diets,
dispersal, 561
phenotypic plasticity, 561
reproductive capacity, 561
small hive beetle, 561
diffusion model,
behavior, 823
dispersal, 823
habitat selection, 823
perceptual range, 823
5,17-dimethylnonacosane,
cuticular hydrocarbons, 1267
9-heptacosyne, 1267
mating behavior, 1267
9-pentacosyne, 1267
Diorhabda elongata,
host choice, 1373
introduced species, 1373
larval performance, 1373
tamarisk biocontrol, 1373
Diptera,
forest management, 657
intensive forestry, 657
pine plantation, 657
water sources, 657
directional orientation,
dispersal, 433
edge effects, 433
immigration, 433
pheromone control, 433
disease management,
epidemiology, 345
vector biology, 345
disease-pathogen dynamics,
Aeromonas salmonicida, 1022
insect feeding assay, 1022
Plecoptera, 1022
trophic interactions, 1022
dispersal, 176, 433, 561, 677, 823, 1005, 1145, 1312
behavior, 823
biodiversity, 1005
biological control, 677
Bt corn, 176
coarse woody debris, 1005

colonization, 1005
diets, 561
diffusion model, 823
directional orientation, 433
edge effects, 433
epigaeic, 1005
European corn borer, 677
gene flow, 1312
habitat selection, 823
herbivory, 1145
immigration, 433
insect resistance management, 176
integrated pest management, 677
invasive species, 1145
microsatellites, 1312
Ostrinia nubilalis, 1312
perceptual range, 823
phenotypic plasticity, 561
pheromone control, 433
population genetics, 1312
postrelease evaluation, 1145
reproductive capacity, 561
small hive beetle, 561
Trichogramma, 677
weed biological control, 1145
western corn rootworm, 176
dispersal behavior,
Diaphorina citri, 1250
huanglongbing, 1250
immunomarking, 1250
mark-capture, 1250
dispersal potential,
age, 235
flight mill, 235
photoperiod, 235
ragweed beetle, 235
distribution, 1038, 1069
bacterial symbionts, 1069
brown stink bug, 1038
Frankliniella occidentalis, 1069
prevalence, 1069
raster map, 1038
southern green stink bug, 1038
distributional pattern,
canonical correlation analysis, 1600
canopy trap, 1600
elevational distribution, 1600
transect, 1600
diversity, 608, 1053
Arribes del Duero Natural Park, 608
biogeography, 608
Carabidae, 1053
ecotones between regenerating and mature stands, 1053
edge-effects, 1053
Mediterranean region, 608
solitary wasps, 608
diving beetle,
Anura, 856
fish, 856
Odonata, 856
predator, 856
dogwood borer,
apple, 1219
behavior, 1219
Sesiidae, 1219
Synanthedon scitula, 1219
dominance,
ant community, 317
citrus orchard, 317
Mediterranean region, 317
seasonality, 317
dryland salinity,
arbovirus, 1585
community, 1585
secondary salinization, 1585
temporary water, 1585
dung beetles,
insect growth regulator, 493
methoprene, 493
Onthophagus taurus, 493

E

East African stemborer species,
host acceptability and suitability, 904
life table, 904
Telenomus isis, 904
temperature and humidity requirements, 904
eastern hemlock,
canopy arthropods, 53
hemlock woolly adelgid, 53
horticultural oil, 53
imidacloprid, 53
ecological disturbance,
conservation management, 325
mediterranean ecosystems, 325
restoration, 325
ecological risk assessment,
biological control of weeds, 731
Cirsium, 731
Trichosirocalus horridus, 731
weed control, 731
ecology,
host fruit, 116
microorganisms, 116
Tephritidae, 116
trapping, 116
economic threshold,
Anaphothrips obscurus, 1737
hybrid economic injury level, 1737
multilinear regression, 1737
ecotones between regenerating and mature stands,
Carabidae, 1053
diversity, 1053
edge-effects, 1053
ectomycorrhiza,
arbuscular mycorrhiza, 93
insect herbivory, 93
insect pollination, 93
edge effects, 43, 433, 1053
Carabidae, 1053
directional orientation, 433
dispersal, 433
diversity, 1053
ecotones between regenerating and mature stands, 1053
fragmentation, 43
immigration, 433
nonmetric multidimensional scaling, 43
pheromone control, 433
sacred groves, 43
West Africa, 43
egg,
ovary, 955
sweet cherry, 955
tart cherry, 955
egg maturation,
Bactrocera oleae, 1496
high temperature, 1496
reproduction, 1496
survival, 1496
egg phenology,
age-dependent developmental rate, 18
phenology, 18
temperature, 18
egg size,
C. transversoguttata, 1283
Coccinella septempunctata, 1283
egg viability, 1283
reproduction, 1283
egg viability,
C. transversoguttata, 1283
Coccinella septempunctata, 1283
egg size, 1283
reproduction, 1283
electroantennogram, 861, 870
attractant, 870
gas chromatographic-electroantennographic detection, 870
gas chromatography-mass spectrometry, 861
headspace analysis, 861
highbush blueberry, 861
pheromone, 870
weevil, 870
Y-tube bioassay, 861
elevated CO_2,
competition, 26
Rhopalosiphum padi, 26
Schizaphis graminum, 26

Sitobion avenae, 26
elevational distribution,
canonical correlation analysis, 1600
canopy trap, 1600
distributional pattern, 1600
transect, 1600
elevational gradient,
herbivory, 1161
phenolics, 1161
plant quality, 1161
emerald ash borer,
Agrilus planipennis, 1226
Fraxinus, 1226
trap tree, 1226
tree stress, 1226
emigration,
aphid behavior, 1429
Myzus persicae, 1429
potato, 1429
virus-induced volatiles, 1429
endophyte, 417, 723
biological control, 723
community interactions, 417
gypsy moth, 417
lesser chestnut weevil, 417
poppy stem gall wasp, 723
systemic protection, 723
Timaspis papaveris, 723
Torymus, 417
endosymbiont,
life table parameters, 1365
psocids, 1365
thermal treatment, 1365
endured resistance,
fall armyworm, 1174
jasmonic acid, 1174
Oryza sativa, 1174
rice, 1174
Spodoptera frugiperda, 1174
Engelmann spruce weevil,
angiosperm volatiles, 1189
nonhost volatiles, 1189
plant–insect interactions, 1189
spatial point processes, 1189
environmental conditions,
density dependent, 962
fertilization, 962
plant morphology, 962
reproduction, 962
environmental factors,
Medicago sativa L., 1480
pollinator visitation, 1480
wild bee, 1480
Epichloë,
context-dependent interactions, 1086
food choice, 1086
Neotyphodium, 1086
Spodoptera frugiperda, 1086
epidemiology,
disease management, 345
vector biology, 345
epigaeic,
biodiversity, 1005
coarse woody debris, 1005
colonization, 1005
dispersal, 1005
ergot alkaloids,
Neotyphodium, 996
tall fescue, 996
trophic interactions, 996
Erwinia tracheiphila,
Acalymma vittatum, 395
frass, 395
polymerase chain reaction, 395
vector–pathogen interactions, 395
Escovopsis,
Attini, 78
Leucoagaricus, 78
Pseudonocardia, 78
symbiosis, 78
ethologic control,
integrated pest management, 781
olfactometric behavior, 781
weevil, 781
ethological software,
adhesives, 168
behavior, 168
bonding, 168
cyanoacrylate, 168
European corn borer,
biological control, 677
dispersal, 677
integrated pest management, 677
Trichogramma, 677
Euschistus servus,
Acrosternum hilare, 450
Glycine max, 450
Nezara viridula, 450
trap crop, 450
exotic insect,
beetle, 266
bidirectional selection, 266
realized heritability, 266
repeatability, 266

F

Fagus grandifolia,
beech bark disease, 1235
beech scale, 1235
Cryptococcus fagisuga, 1235
fall armyworm,
endured resistance, 1174
jasmonic acid, 1174
Oryza sativa, 1174
rice, 1174
Spodoptera frugiperda, 1174
fecundity,
development, 1661
oviposition, 1661
Solanaceae, 1661
survival, 1661
feeding,
Cicadellidae, 1485
glassy-winged sharpshooter, 1485
survival, 1485
temperature, 1485
vector, 1485
feeding behavior,
biological control, 715
phytophagous mites, 715
predator–prey interaction, 715
predatory potential, 715
fertility,
low temperature, 242
males, 242
mating, 242
mobility, 242
parasitoids, 242
fertilization,
density dependent, 962
environmental conditions, 962
plant morphology, 962
reproduction, 962
field cage,
Chrysopa nigricornis, 1117
field observation, 1117
Harmonia axyridis, 1117
laboratory arena, 1117
field monitoring,
Bt corn, 281
Cry1F, 281
nontarget arthropods, 281
field observation,
Chrysopa nigricornis, 1117
field cage, 1117
Harmonia axyridis, 1117
laboratory arena, 1117
field survey,
Apis mellifera, 514
colony health, 514
France, 514
pesticides, 514
fire ants,
Argentina, 198
biological control, 198
parasitoid, 198
social form, 198
firefly,
climate change, 1505
degree-day, 1505
phenology, 1505
synchronous display, 1505
fish,
Anura, 856
diving beetle, 856
Odonata, 856
predator, 856
fitness,
isogenic strain, 499
M factor, 499
sex determination, 499

flight activity,
Acrosternum hilare, 1716
black light trap, 1716
degree-day, 1716
population dynamics, 1716
flight behavior,
automated trapping, 1555
Lymantria dispar, 1555
phenology, 1555
sampling, 1555
flight mill,
age, 235
dispersal potential, 235
photoperiod, 235
ragweed beetle, 235
flight phenology,
chronological reproductive isolation, 1697
Macrotermes, 1697
swarming, 1697
Florida,
host plant apparency, 1211
larval performance, 1211
oviposition preference, 1211
Yucca filamentosa, 1211
flower density, 748, 1680
Bombus appositus, 1680
bumble bee, 748, 1680
Delphinium barbeyi, 1680
food availability, 748
food limitation, 1680
foraging distance, 748
subalpine, 748
fly production,
ammonia, 633
biosolid cake, 633
temperature, 633
food availability,
bumble bee, 748
flower density, 748
foraging distance, 748
subalpine, 748
food chain,
heavy metals, 1276
lead, 1276
mulberry, 1276
silkworm, 1276
food choice,
context-dependent interactions, 1086
Epichloë, 1086
Neotyphodium, 1086
Spodoptera frugiperda, 1086
food limitation,
Bombus appositus, 1680
bumble bee, 1680
Delphinium barbeyi, 1680
flower density, 1680
foraging, 1241, 1379, 1618
allocation, 1379
ant–aphid interaction, 1618
Coptotermes formosanus, 1241
Helicoverpa zea, 1379
integrated pest management, 1241
nesting, 1241
nutrition, 1379
precipitation, 1618
Solenopsis invicta, 1241
starvation, 1379
foraging behavior,
marginal value theorem, 183
spatial analysis of distance indices, 183
foraging distance,
bumble bee, 748
flower density, 748
food availability, 748
subalpine, 748
forest,
apple, 530
ornamentals, 530
pheromone trapping, 530
Synanthedon scitula, 530
forest management,
Diptera, 657
intensive forestry, 657
pine plantation, 657
water sources, 657
Formicidae,
inventory, 1724
Ohio, 1724
sampling methods, 1724
species richness, 1724
Fragaria spp.,
precision pest management, 137
spider mites, 137
visible and near infrared spectra, 137
fragmentation,
edge effects, 43
nonmetric multidimensional scaling, 43
sacred groves, 43
West Africa, 43
France,
Apis mellifera, 514
colony health, 514
field survey, 514
pesticides, 514
Frankliniella australis,
Frankliniella occidentalis, 333
Orius, 333
population abundance, 333
predation, 333
Frankliniella fusca,
Frankliniella occidentalis, 1205
Frankliniella tritici, 1205
population dynamics, 1205
within-plant aggregation, 1205
Frankliniella occidentalis, 333, 1069, 1205
bacterial symbionts, 1069
distribution, 1069
Frankliniella australis, 333
Frankliniella fusca, 1205
Frankliniella tritici, 1205
Orius, 333
population abundance, 333
population dynamics, 1205
predation, 333
prevalence, 1069
within-plant aggregation, 1205
Frankliniella tritici,
Frankliniella fusca, 1205
Frankliniella occidentalis, 1205
population dynamics, 1205
within-plant aggregation, 1205
frass,
Acalymma vittatum, 395
Erwinia tracheiphila, 395
polymerase chain reaction, 395
vector–pathogen interactions, 395
Fraxinus,
Agrilus planipennis, 1226
emerald ash borer, 1226
trap tree, 1226
tree stress, 1226
frequency distribution,
codling moth, 1395
Cydia pomonella, 1395
larval aggregation, 1395
pheromone, 1395
frontalin,
bark beetle phermone, 472
cis-verbenol, 472
red turpentine beetle, 472
trans-verbenol, 472
functional response, 708, 1634
biological control, 708, 1634
Gonatocerus ashmeadi, 1634
Gonatocerus tuberculifemur, 1634
Homalodisca vitripennis, 1634
multicolored Asian lady beetle, 708
seven-spotted lady beetle, 708
soybean aphid, 708
fungal transmission,
ophiostomatoid fungi, 143
phoresy, 143
Tarsonemus, 143
vector, 143

G

gall insect,
Aucuba japonica, 404
physical defense, 404
plant–herbivore interaction, 404
susceptible period, 404
gas chromatographic-electroantennographic detection,
attractant, 870
electroantennogram, 870

pheromone, 870
weevil, 870
gas chromatography-mass spectrometry,
electroantennogram, 861
headspace analysis, 861
highbush blueberry, 861
Y-tube bioassay, 861
gene flow,
dispersal, 1312
microsatellites, 1312
Ostrinia nubilalis, 1312
population genetics, 1312
genetic structure,
ambrosia beetle, 1324
cytochrome oxidase I, 1324
nested clade phylogeographic analysis, 1324
genetically modified organisms,
carabid beetles, 944
corn, 944
Orius, 944
spiders, 944
girdled trees,
detection methods, 1668
Manuka oil, 1668
methyl jasmonate, 1668
trap tree, 1668
girdling,
Buprestidae, 1756
Coleoptera, 1756
insect–plant interactions, 1756
light intensity, 1756
glassy-winged sharpshooter,
Cicadellidae, 1485
feeding, 1485
survival, 1485
temperature, 1485
vector, 1485
glycerol,
Chilo suppressalis, 260
cold hardiness, 260
diapause, 260
rice stem borer, 260
Glycine max,
Acrosternum hilare, 450
Euschistus servus, 450
Nezara viridula, 450
trap crop, 450
Gonatocerus ashmeadi,
biological control, 1634
functional response, 1634
Gonatocerus tuberculifemur, 1634
Homalodisca vitripennis, 1634
Gonatocerus tuberculifemur,
biological control, 1634
functional response, 1634
Gonatocerus ashmeadi, 1634
Homalodisca vitripennis, 1634
grasshopper development,
degree-day, 1521
phenology, 1521
rangeland, 1521
grazing pressure,
arachnids, 985
breeds, 985
insects, 985
ungulates, 985
Great Basin,
agroecology, 1028
agroecosystem, 1028
cross-boundary, 1028
spillover, 1028
green peach aphid,
aphid population fitness, 129
Myzus persicae, 129
peroxidase, 129
phenolics, 129
gut microbiota,
Anoplophora glabripennis, 686
culture-independent community analysis, 686
host tree resistance, 686
gypsy moth,
community interactions, 417
endophyte, 417
lesser chestnut weevil, 417
Torymus, 417

H

habitat microenvironment,
Neoseiulus, 920
solar UV radiation, 920
Tetranychus, 920
habitat selection,
behavior, 823
diffusion model, 823
dispersal, 823
perceptual range, 823
haplotype,
cytochrome oxidase, 1546
Noctuidae, 1546
Spodoptera frugiperda, 1546
Harmonia axyridis, 1117, 1563
Chrysopa nigricornis, 1117
contaminant pest, 1563
field cage, 1117
field observation, 1117
laboratory arena, 1117
modeling, 1563
sugar content, 1563
trap catching, 1563
Harpalus pensylvanicus,
Bt, 935
Carabidae, 935
Cry3Bb, 935
neonicotinoids, 935
headspace analysis,
electroantennogram, 861
gas chromatography-mass spectrometry, 861
highbush blueberry, 861
Y-tube bioassay, 861
heavy metals,
food chain, 1276
lead, 1276
mulberry, 1276
silkworm, 1276
Helicoverpa armigera,
Bt cotton, 1336
Cry1Ac, 1336
resistance frequency, 1336
Helicoverpa zea,
allocation, 1379
foraging, 1379
nutrition, 1379
starvation, 1379
hemlock,
Adelges tsugae, 577
climate, 577
invasive species, 577
landscape, 577
hemlock woolly adelgid,
canopy arthropods, 53
eastern hemlock, 53
horticultural oil, 53
imidacloprid, 53
9-heptacosyne,
cuticular hydrocarbons, 1267
5,17-dimethylnonacosane, 1267
mating behavior, 1267
9-pentacosyne, 1267
herbivore guild,
physiological structure, 667
plant–insect dynamics, 667
productivity, 667
resource competition, 667
herbivory, 1145, 1161
dispersal, 1145
elevational gradient, 1161
invasive species, 1145
phenolics, 1161
plant quality, 1161
postrelease evaluation, 1145
weed biological control, 1145
Hermetia illucens,
black soldier fly, 930
development, 930
temperature, 930
Heteroptera,
Buchnera, 103
southern green stink bug, 103
symbiosis, 103
Heterorhabditis,
Beauveria bassiana, 1126
inundative biological control, 1126
Metarhizium anisopliae, 1126
Steinernema, 1126
high temperature,
Bactrocera oleae, 1496
egg maturation, 1496
reproduction, 1496
survival, 1496
highbush blueberry,
electroantennogram, 861
gas chromatography-mass spectrometry, 861
headspace analysis, 861
Y-tube bioassay, 861
Homalodisca vitripennis,
biological control, 1634
functional response, 1634

Gonatocerus ashmeadi, 1634
Gonatocerus tuberculifemur, 1634
Hoplitis,
Megachile, 1293
Osmia, 1293
pollinator performance, 1293
resource limitation, 1293
horned passalus,
bess beetle, 1411
coarse woody debris, 1411
landscape matrix, 1411
horticultural oil,
canopy arthropods, 53
eastern hemlock, 53
hemlock woolly adelgid, 53
imidacloprid, 53
host acceptability and suitability,
East African stemborer species, 904
life table, 904
Telenomus isis, 904
temperature and humidity requirements, 904
host age,
host density, 1513
Lobesia botrana, 1513
Trichogramma cacoeciae, 1513
UV treatment, 1513
host choice,
Diorhabda elongata, 1373
introduced species, 1373
larval performance, 1373
tamarisk biocontrol, 1373
host density,
host age, 1513
Lobesia botrana, 1513
Trichogramma cacoeciae, 1513
UV treatment, 1513
host finding,
aphid–plant interaction, 411
Carya illinoinensis, 411
leaf chlorosis, 411
Melanocallis caryaefoliae, 411
host fruit,
ecology, 116
microorganisms, 116
Tephritidae, 116
trapping, 116
host plant,
cotton fleahopper, 766
cotton pest, 766
host plant apparency,
Florida, 1211
larval performance, 1211
oviposition preference, 1211
Yucca filamentosa, 1211
host plant volatiles,
Anthonomus eugenii, 216
Capsicum annuum, 216
Solanum americanum, 216
Solanum melongena, 216
host preference,
bark beetles, 1197
Dendroctonus rhizophagus, 1197
pine seedlings, 1197
tree mortality, 1197
host preference and performance,
insect–plant interaction, 1653
invasion, 1653
Nuphar lutea, 1653
Trapa natans, 1653
host selection, 797, 896, 1707
biological control, 896
Cerambycidae, 797
circadian rhythm, 896
climatic parameters, 896
insect–plant interactions, 1707
intraspecific competition, 1707
oviposition behavior, 1707
physiological state, 896
primary attraction, 797
Scolytinae, 797
semiochemicals, 797
host specificity,
biological weed control, 1135
China, 1135
invasive plant, 1135
leaf-rolling weevil, 1135
host suitability,
crapemyrtle, 1155
crapemyrtle aphid, 1155
host tree resistance,
Anoplophora glabripennis, 686
culture-independent community analysis, 686
gut microbiota, 686
host volatiles,
native versus exotic hosts, 757
olfactory cue, 757
host-parasitoid,
density dependence, 790
Identificaticoffee agroecosystem, 790
olfactory cue, 790
scale-dependent effects, 790
host–parasitoid interaction,
biological control, 365
host–parasitoid interaction, 365
Lycopersicon esculentum, 365
spatial scales, 365
house fly,
Muscidifurax raptorellus, 700
stable fly, 700
Tachinaephagus zealandicus, 700
Trichopria nigra, 700
huanglongbing,
Diaphorina citri, 1250
dispersal behavior, 1250
immunomarking, 1250
mark-capture, 1250
hybrid economic injury level,
Anaphothrips obscurus, 1737
economic threshold, 1737
multilinear regression, 1737

I

Identificaticoffee agroecosystem,
density dependence, 790
host-parasitoid, 790
olfactory cue, 790
scale-dependent effects, 790
imidacloprid,
canopy arthropods, 53
eastern hemlock, 53
hemlock woolly adelgid, 53
horticultural oil, 53
immigration,
directional orientation, 433
dispersal, 433
edge effects, 433
pheromone control, 433
immunomarking,
Diaphorina citri, 1250
dispersal behavior, 1250
huanglongbing, 1250
mark-capture, 1250
in situ enclosures,
instantaneous growth rate, 250
surface-floating pupal exuviae, 250
winter active, 250
insect agriculture,
artificial medium, 1096
biological invasion, 1096
cooperative breeding, 1096
symbiosis, 1096
insect demography,
biological control, 2
multiple decrement life table, 2
natural enemies, 2
risk analysis, 2
insect feeding assay,
Aeromonas salmonicida, 1022
disease-pathogen dynamics, 1022
Plecoptera, 1022
trophic interactions, 1022
insect growth regulator,
dung beetles, 493
methoprene, 493
Onthophagus taurus, 493
insect herbivory,
arbuscular mycorrhiza, 93
ectomycorrhiza, 93
insect pollination, 93
insect pathogens,
Cephus cinctus, 387
plant pathogens, 387
Triticum aestivum, 387
wheat, 387
insect pollination,
arbuscular mycorrhiza, 93
ectomycorrhiza, 93
insect herbivory, 93

insect resistance management,
Bt corn, 176
dispersal, 176
western corn rootworm, 176
insect symbiont,
bacterial interaction, 1360
biological control, 1360
insect/fungus associations,
ophiostomatoid fungi, 356
tree disease, 356
insect–plant interactions, 1653, 1707, 1756
Buprestidae, 1756
Coleoptera, 1756
girdling, 1756
host preference and performance, 1653
host selection, 1707
intraspecific competition, 1707
invasion, 1653
light intensity, 1756
Nuphar lutea, 1653
oviposition behavior, 1707
Trapa natans, 1653
insects, 67, 985
arachnids, 985
bark beetles, 67
breeds, 985
context dependency, 67
grazing pressure, 985
microorganisms, 67
symbiosis, 67
ungulates, 985
instantaneous growth rate,
in situ enclosures, 250
surface-floating pupal exuviae, 250
winter active, 250
integrated pest management, 677, 781, 809, 1106, 1168, 1241
biological control, 677
Coptotermes formosanus, 1241
Cylindroiulus caeruleocinctus, 1106
Diabrotica undecimpunctata undecimpuncta, 809
dispersal, 677
ethologic control, 781
European corn borer, 677
foraging, 1241
nesting, 1241
olfactometric behavior, 781
Pterostichus melanarius, 1106
ruthenium red dye, 1168
sampling plan design, 809
scouting programs, 809
Solenopsis invicta, 1241
southern green stink bug, 1168
strip spraying, 809
Trichogramma, 677
weevil, 781
intensive forestry,
Diptera, 657
forest management, 657
pine plantation, 657
water sources, 657
intercropping,
Bemisia argentifolii, 442
living mulch, 442
phytotoxemia, 442
squash, 442
interspecific competition,
aggression, 425
longhorned beetle, 425
mating behavior, 425
intraspecific competition,
host selection, 1707
insect–plant interactions, 1707
oviposition behavior, 1707
intrinsic rate of increase,
development, 1061
meta-analysis, 1061
reproduction, 1061
Russian wheat aphid, 1061
introduced species,
Diorhabda elongata, 1373
host choice, 1373
larval performance, 1373
tamarisk biocontrol, 1373
inundative biological control,
Beauveria bassiana, 1126
Heterorhabdtis, 1126
Metarhizium anisopliae, 1126
Steinernema, 1126
invasion,
host preference and performance, 1653
insect–plant interaction, 1653
Nuphar lutea, 1653
Trapa natans, 1653
invasive plant,
biological weed control, 1135
China, 1135
host specificity, 1135
leaf-rolling weevil, 1135
invasive species, 577, 1145, 1400
Adelges tsugae, 577
climate, 577
dispersal, 1145
hemlock, 577
herbivory, 1145
landscape, 577
larval behavior, 1400
Olea europea, 1400
population dispersion, 1400
postrelease evaluation, 1145
weed biological control, 1145
inventory,
Formicidae, 1724
Ohio, 1724
sampling methods, 1724
species richness, 1724
isogenic strain,
fitness, 499
M factor, 499
sex determination, 499

J

Japanese barberry,
blacklegged tick, 977
Lyme disease, 977
white-footed mouse, 977
white-tailed deer, 977
jasmonic acid,
endured resistance, 1174
fall armyworm, 1174
Oryza sativa, 1174
rice, 1174
Spodoptera frugiperda, 1174

K

kairomones,
Asian citrus psyllid, 225
attractant, 225
citrus greening disease, 225
olfaction, 225

L

laboratory arena,
Chrysopa nigricornis, 1117
field cage, 1117
field observation, 1117
Harmonia axyridis, 1117
laboratory experiments,
biosafety, 293
natural enemies, 293
review, 293
transgenic plants, 293
landscape,
Adelges tsugae, 577
climate, 577
hemlock, 577
invasive species, 577
landscape ecology,
remote sensing, 551
Solenopsis invicta, 551
landscape matrix,
bess beetle, 1411
coarse woody debris, 1411
horned passalus, 1411
Laodelphax striatellus,
cytoplasmic incompatibility, 1626
microinjection, 1626
Nilaparvata lugens, 1626
Wolbachia, 1626
larvae,
baiting, 35
castes, 35
reproductive biology, 35
subterranean termites, 35
larval aggregation,
codling moth, 1395
Cydia pomonella, 1395
frequency distribution, 1395
pheromone, 1395
larval behavior,
invasive species, 1400
Olea europea, 1400
population dispersion, 1400

larval development,
Agapeta zoegana, 971
biological control, 971
Centaurea stoebe, 971
Cyphocleonus achates, 971
diapause, 971
larval habitats,
Aedes vigilax, 1013
Culex annulirostris, 1013
density dependence, 1013
vector control, 1013
larval performance, 1211, 1373
Diorhabda elongata, 1373
Florida, 1211
host choice, 1373
host plant apparency, 1211
introduced species, 1373
oviposition preference, 1211
tamarisk biocontrol, 1373
Yucca filamentosa, 1211
lead,
food chain, 1276
heavy metals, 1276
mulberry, 1276
silkworm, 1276
leaf chlorosis,
aphid–plant interaction, 411
Carya illinoinensis, 411
host finding, 411
Melanocallis caryaefoliae, 411
leaf-rolling weevil,
biological weed control, 1135
China, 1135
host specificity, 1135
invasive plant, 1135
learning,
memory, 803
plasticity, 803
social wasps, 803
lesser chestnut weevil,
community interactions, 417
endophyte, 417
gypsy moth, 417
Torymus, 417
Leucoagaricus,
Attini, 78
Escovopsis, 78
Pseudonocardia, 78
symbiosis, 78
life cycle,
developmental period, 478
powderpost beetle, 478
thermal constants, 478
thermal requirements, 478
life table,
East African stemborer species, 904
host acceptability and suitability, 904
Telenomus isis, 904
temperature and humidity requirements, 904
life table parameters,
endosymbiont, 1365
psocids, 1365
thermal treatment, 1365
light intensity,
Buprestidae, 1756
Coleoptera, 1756
girdling, 1756
insect–plant interactions, 1756
linear and nonlinear models,
codling moth, 885
Cydia pomonella, 885
development, 885
thermal threshold, 885
living mulch,
Bemisia argentifolii, 442
intercropping, 442
phytotoxemia, 442
squash, 442
Lobesia botrana,
host age, 1513
host density, 1513
Trichogramma cacoeciae, 1513
UV treatment, 1513
locust,
diapause, 569
Madagascar, 569
reproductive maturity, 569
wing color, 569
longhorned beetle, 425, 1462
aggression, 425
interspecific competition, 425
mating behavior, 425, 1462
Mojave desert, 1462
(*R*)-3-hydroxyhexan-2-one, 1462
sex pheromone, 1462
low temperature, 242, 1539
Aphthona flea beetles, 1539
chill degree-days, 1539
fertility, 242
males, 242
mating, 242
mobility, 242
overwintering, 1539
parasitoid, 242
snow cover, 1539
Lycopersicon esculentum,
biological control, 365
host–parasitoid interaction, 365
spatial scales, 365
Lymantria dispar,
automated trapping, 1555
flight behavior, 1555
phenology, 1555
sampling, 1555
Lyme disease,
blacklegged tick, 977
Japanese barberry, 977
white-footed mouse, 977
white-tailed deer, 977

M

M factor,
fitness, 499
isogenic strain, 499
sex determination, 499
Macrotermes,
chronological reproductive isolation, 1697
flight phenology, 1697
swarming, 1697
Madagascar,
diapause, 569
locust, 569
reproductive maturity, 569
wing color, 569
Malacosoma disstria,
Choristoneura conflictana, 459
monitor, 459
prediction, 459
sex pheromone traps, 459
male-produced pheromone,
Anoplophora glabripennis, 1745
Cerambycidae, 1745
mate-finding, 1745
trap design, 1745
males,
fertility, 242
low temperature, 242
mating, 242
mobility, 242
parasitoid, 242
Mangora gibberosa, 651, 1595
Araneidae, 651
Argiope trifasciata, 651, 1595
Cyclosa turbinata, 651, 1595
Neoscona arabesca, 651, 1595
species diversity, 1595
Manuka oil,
detection methods, 1668
girdled trees, 1668
methyl jasmonate, 1668
trap tree, 1668
marginal value theorem,
foraging behavior, 183
spatial analysis of distance indices, 183
mark-capture,
Diaphorina citri, 1250
dispersal behavior, 1250
huanglongbing, 1250
immunomarking, 1250
mate-finding,
Anoplophora glabripennis, 1745
Cerambycidae, 1745
male-produced pheromone, 1745
trap design, 1745
mating,
fertility, 242
low temperature, 242
males, 242
mobility, 242
parasitoid, 242
mating behavior, 192, 425, 1267, 1462
Aegorhinus superciliosus, 192
aggression, 425
contact pheromone, 192
cuticular hydrocarbons, 1267
5,17-dimethylnonacosane, 1267

9-heptacosyne, 1267
interspecific competition, 425
longhorned beetle, 425, 1462
Mojave desert, 1462
9-pentacosyne, 1267
(*R*)-3-hydroxyhexan-2-one, 1462
raspberry weevil, 192
sex pheromone, 1462
Medicago sativa L.,
environmental factors, 1480
pollinator visitation, 1480
wild bee, 1480
mediterranean ecosystems,
conservation management, 325
ecological disturbance, 325
restoration, 325
Mediterranean region, 317, 608
ant community, 317
Arribes del Duero Natural Park, 608
biogeography, 608
citrus orchard, 317
diversity, 608
dominance, 317
seasonality, 317
solitary wasps, 608
Megachile,
Hoplitis, 1293
Osmia, 1293
pollinator performance, 1293
resource limitation, 1293
Melanocallis caryaefoliae,
aphid–plant interaction, 411
Carya illinoinensis, 411
host finding, 411
leaf chlorosis, 411
memory,
learning, 803
plasticity, 803
social wasps, 803
meta-analysis,
development, 1061
intrinsic rate of increase, 1061
reproduction, 1061
Russian wheat aphid, 1061
Metarhizium anisopliae,
Beauveria bassiana, 1126
Heterorhabdtis, 1126
inundative biological control, 1126
Steinernema, 1126
methoprene,
dung beetles, 493
insect growth regulator, 493
Onthophagus taurus, 493
methyl eugenol,
Bactrocera, 1446
Cue-Lure, 1446
methyl jasmonate, 1439, 1668
Ceroplastes japonicus Green, 1439
Chilocorus kuwanae Silvestri, 1439
detection methods, 1668
girdled trees, 1668
Manuka oil, 1668
α-pinene, 1439
trap tree, 1668
volatiles, 1439
2-methyl-4-octanone,
agave, 1423
pheromone release, 1423
weevil, 1423
microinjection,
cytoplasmic incompatibility, 1626
Laodelphax striatellus, 1626
Nilaparvata lugens, 1626
Wolbachia, 1626
microorganisms, 67, 116
bark beetles, 67
context dependency, 67
ecology, 116
host fruit, 116
insects, 67
symbiosis, 67
Tephritidae, 116
trapping, 116
microsatellites, 1301, 1312
Aphis glycines, 1301
dispersal, 1312
gene flow, 1312
Ostrinia nubilalis, 1312
population genetic structure, 1301
population genetics, 1312
soybean aphid, 1301
mobility,
fertility, 242
low temperature, 242
males, 242
mating, 242
parasitoid, 242
modeling,
contaminant pest, 1563
Harmonia axyridis, 1563
sugar content, 1563
trap catching, 1563
Mojave desert,
longhorned beetle, 1462
mating behavior, 1462
(*R*)-3-hydroxyhexan-2-one, 1462
sex pheromone, 1462
monitor,
Choristoneura conflictana, 459
Malacosoma disstria, 459
prediction, 459
sex pheromone traps, 459
mulberry,
food chain, 1276
heavy metals, 1276
lead, 1276
silkworm, 1276
multicolored Asian lady beetle,
biological control, 708
functional response, 708
seven-spotted lady beetle, 708
soybean aphid, 708
multilinear regression,
Anaphothrips obscurus, 1737
economic threshold, 1737
hybrid economic injury level, 1737
multiple decrement life table,
biological control, 2
insect demography, 2
natural enemies, 2
risk analysis, 2
multispecies interactions,
Apis mellifera, 207
Bombus spp., 207
Peponapis pruinosa, 207
winter squash, 207
Muscidifurax raptorellus,
house fly, 700
stable fly, 700
Tachinaephagus zealandicus, 700
Trichopria nigra, 700
mutualism,
antagonism, 741
Apocynaceae, 741
nectar, 741
pollination syndromes, 741
Myzus persicae, 129, 153, 1429
aphid behavior, 1429
aphid population fitness, 129
Aphidius colemani, 153
aphids, 153
Coccinella undecimpunctata, 153
emigration, 1429
green peach aphid, 129
peroxidase, 129
phenolics, 129
potato, 1429
prey quality, 153
virus-induced volatiles, 1429

N

NADH dehydrogenase 1,
biological invasion, 274
Striacosta albicosta, 274
Zea mays, 274
native,
ant, 846
coccinellid, 846
natural enemies, 846
non-native, 846
native versus exotic hosts,
host volatiles, 757
olfactory cue, 757
natural enemies, 2, 159, 293, 846, 1642
ant, 846
biological control, 2
biosafety, 293
Brazilian peppertree, 1642
Christmas berry, 1642
coccinellid, 846
insect demography, 2
laboratory experiments, 293

multiple decrement life table, 2
native, 846
non-native, 846
olfactometer, 159
phenylacetaldehyde, 159
review, 293
risk analysis, 2
risk assessment, 1642
transgenic plants, 293
visual cues, 159
western tarnished plant bug, 159
nectar,
antagonism, 741
Apocynaceae, 741
mutualism, 741
pollination syndromes, 741
neonicotinoids,
Bt, 935
Carabidae, 935
Cry3Bb, 935
Harpalus pensylvanicus, 935
Neoscona arabesca, 651, 1595
Araneidae, 651
Argiope trifasciata, 651, 1595
Cyclosa turbinata, 651, 1595
Mangora gibberosa, 651, 1595
species diversity, 1595
Neoseiulus,
habitat microenvironment, 920
solar UV radiation, 920
Tetranychus, 920
Neotyphodium, 996, 1086
context-dependent interactions, 1086
Epichloë, 1086
ergot alkaloids, 996
food choice, 1086
Spodoptera frugiperda, 1086
tall fescue, 996
trophic interactions, 996
nest site selection,
biological invasion, 484
parasitoid, 484
stress, 484
temperature, 484
nest-site limitation,
ant nest, 600
urban ecology, 600
nested clade phylogeographic analysis,
ambrosia beetle, 1324
cytochrome oxidase I, 1324
genetic structure, 1324
nesting,
Coptotermes formosanus, 1241
foraging, 1241
integrated pest management, 1241
Solenopsis invicta, 1241
Nezara viridula,
Acrosternum hilare, 450
Euschistus servus, 450
Glycine max, 450
trap crop, 450
Nilaparvata lugens,
cytoplasmic incompatibility, 1626
Laodelphax striatellus, 1626
microinjection, 1626
Wolbachia, 1626
Noctuidae,
cytochrome oxidase, 1546
haplotype, 1546
Spodoptera frugiperda, 1546
non-native,
ant, 846
coccinellid, 846
native, 846
natural enemies, 846
nonhost volatiles,
angiosperm volatiles, 1189
Engelmann spruce weevil, 1189
plant–insect interactions, 1189
spatial point processes, 1189
nonmetric multidimensional scaling,
edge effects, 43
fragmentation, 43
sacred groves, 43
West Africa, 43
nontarget arthropods,
Bt corn, 281
Cry1F, 281
field monitoring, 281
North America,
Braconidae, 588
co-evolution, 588
Rhagoletis pomonella, 588
Tephritidae, 588
Nuphar lutea,
host preference and performance, 1653
insect–plant interaction, 1653
invasion, 1653
Trapa natans, 1653
nutrition,
allocation, 1379
foraging, 1379
Helicoverpa zea, 1379
starvation, 1379
nymphal survival,
Brassica oleracea L. variety *italica* Plenck, 11
development, 11
phenological models, 11

O

Odonata,
Anura, 856
diving beetle, 856
fish, 856
predator, 856
Ohio,
Formicidae, 1724
inventory, 1724
sampling methods, 1724
species richness, 1724
Olea europea,
invasive species, 1400
larval behavior, 1400
population dispersion, 1400
olfaction, 225, 1259
Asian citrus psyllid, 225
attractant, 225
Caribbean fruit fly, 1259
citrus greening disease, 225
1,4-diaminobutane, 1259
kairomones, 225
semiochemicals, 1259
synthetic attractants, 1259
olfactometer,
natural enemies, 159
phenylacetaldehyde, 159
visual cues, 159
western tarnished plant bug, 159
olfactometer bioassay,
crucifer flea beetle, 1467
volatile compounds, 1467
olfactometric behavior,
ethologic control, 781
integrated pest management, 781
weevil, 781
olfactory cue, 757, 790
density dependence, 790
host volatiles, 757
host-parasitoid, 790
Identificaticoffee agroecosystem, 790
native versus exotic hosts, 757
scale-dependent effects, 790
onion,
colonization process, 1575
population model, 1575
Thrips tabaci, 1575
Onthophagus taurus,
dung beetles, 493
insect growth regulator, 493
methoprene, 493
ophiostomatoid fungi, 143, 356
fungal transmission, 143
insect/fungus associations, 356
phoresy, 143
Tarsonemus, 143
tree disease, 356
vector, 143
Orius, 333, 944
carabid beetles, 944
corn, 944
Frankliniella australis, 333
Frankliniella occidentalis, 333
genetically modified organisms, 944
population abundance, 333
predation, 333
spiders, 944
ornamentals,
apple, 530
forest, 530
pheromone trapping, 530
Synanthedon scitula, 530

Oryza sativa,
endured resistance, 1174
fall armyworm, 1174
jasmonic acid, 1174
rice, 1174
Spodoptera frugiperda, 1174
Osmia,
Hoplitis, 1293
Megachile, 1293
pollinator performance, 1293
resource limitation, 1293
Ostrinia nubilalis,
dispersal, 1312
gene flow, 1312
microsatellites, 1312
population genetics, 1312
ovary,
egg, 955
sweet cherry, 955
tart cherry, 955
overwintering, 524, 1182, 1539
Aphthona flea beetles, 1539
chill degree-days, 1539
Chilo suppressalis, 1182
cold tolerance, 524
low temperature, 1539
paddy rice, 1182
pest management, 524
Plutella xylostella, 524
snow cover, 1539
supercooling capacity, 1182
water-oat, 1182
oviposition,
development, 1661
fecundity, 1661
Solanaceae, 1661
survival, 1661
oviposition behavior,
host selection, 1707
insect–plant interactions, 1707
intraspecific competition, 1707
oviposition preference,
Florida, 1211
host plant apparency, 1211
larval performance, 1211
Yucca filamentosa, 1211

P

paddy rice,
Chilo suppressalis, 1182
overwintering, 1182
supercooling capacity, 1182
water-oat, 1182
parasitoid, 198, 242, 375, 484
Argentina, 198
biological control, 198
biological invasion, 484
fertility, 242
fire ants, 198
low temperature, 242
males, 242
mating, 242
mobility, 242
nest site selection, 484
Pentatomidae, 375
Scelionidae, 375
social form, 198
stress, 484
Telenomus podisi, 375
temperature, 484
Trissolcus basalis, 375
pea leafminer,
Agral 90, 1387
antixenosis, 1387
preoviposition behaviors, 1387
Sylgard 309, 1387
pear psylla,
Cacopsylla pyricola, 815
diel periodicity, 815
sex attraction, 815
pecan nut casebearer,
Acrobasis nuxvorella, 1690
circadian, 1690
periodicity, 1690
pheromone, 1690
9-pentacosyne,
cuticular hydrocarbons, 1267
5,17-dimethylnonacosane, 1267
9-heptacosyne, 1267
mating behavior, 1267
Pentatomidae,
parasitoid, 375
Scelionidae, 375
Telenomus podisi, 375
Trissolcus basalis, 375
Peponapis pruinosa,
Apis mellifera, 207
Bombus spp., 207
multispecies interactions, 207
winter squash, 207
perceptual range,
behavior, 823
diffusion model, 823
dispersal, 823
habitat selection, 823
periodicity,
Acrobasis nuxvorella, 1690
circadian, 1690
pecan nut casebearer, 1690
pheromone, 1690
peroxidase,
aphid population fitness, 129
green peach aphid, 129
Myzus persicae, 129
phenolics, 129
pest control,
bacteria symbionts, 1078
protease inhibitor, 1078
proteases, 1078
16S rRNA, 1078
pest management,
cold tolerance, 524
overwintering, 524
Plutella xylostella, 524
pesticides,
Apis mellifera, 514
colony health, 514
field survey, 514
France, 514
Phaseolus vulgaris,
Acyrthosiphon pisum, 1347
Aphis glycines, 1347
Aphis gossypii, 1347
Therioaphis trifolii, 1347
phenolics, 129, 1161
aphid population fitness, 129
elevational gradient, 1161
green peach aphid, 129
herbivory, 1161
Myzus persicae, 129
peroxidase, 129
plant quality, 1161
phenological models,
Brassica oleracea L. variety *italica* Plenck, 11
development, 11
nymphal survival, 11
phenology, 18, 307, 539, 1505, 1521, 1555
age-dependent developmental rate, 18
automated trapping, 1555
biological control, 539
climate change, 1505
damage, 307
degree-day, 1505, 1521
egg phenology, 18
firefly, 1505
flight behavior, 1555
grasshopper development, 1521
Lymantria dispar, 1555
precipitation, 307
rangeland, 1521
relative abundance, 539
sampling, 539, 1555
Solenopsis invicta, 539
synchronous display, 1505
temperature, 18, 307
voltinism, 307
phenotypic plasticity,
diets, 561
dispersal, 561
reproductive capacity, 561
small hive beetle, 561
phenylacetaldehyde,
natural enemies, 159
olfactometer, 159
visual cues, 159
western tarnished plant bug, 159
pheromone, 870, 1395, 1690
Acrobasis nuxvorella, 1690
attractant, 870
circadian, 1690
codling moth, 1395
Cydia pomonella, 1395
electroantennogram, 870
frequency distribution, 1395
gas chromatographic-electroantennographic detection, 870
larval aggregation, 1395
pecan nut casebearer, 1690

periodicity, 1690
weevil, 870
pheromone control,
directional orientation, 433
dispersal, 433
edge effects, 433
immigration, 433
pheromone release,
agave, 1423
2-methyl-4-octanone, 1423
weevil, 1423
pheromone trapping,
apple, 530
forest, 530
ornamentals, 530
Synanthedon scitula, 530
phoresy,
fungal transmission, 143
ophiostomatoid fungi, 143
Tarsonemus, 143
vector, 143
photoperiod,
age, 235
dispersal potential, 235
flight mill, 235
ragweed beetle, 235
physical defense,
Aucuba japonica, 404
gall insect, 404
plant–herbivore interaction, 404
susceptible period, 404
physiological state,
biological control, 896
circadian rhythm, 896
climatic parameters, 896
host selection, 896
physiological structure,
herbivore guild, 667
plant–insect dynamics, 667
productivity, 667
resource competition, 667
phytophagous mites,
biological control, 715
feeding behavior, 715
predator–prey interaction, 715
predatory potential, 715
phytotoxemia,
Bemisia argentifolii, 442
intercropping, 442
living mulch, 442
squash, 442
pine plantation,
Diptera, 657
forest management, 657
intensive forestry, 657
water sources, 657
pine seedlings,
bark beetles, 1197
Dendroctonus rhizophagus, 1197
host preference, 1197
tree mortality, 1197
α-pinene,
Ceroplastes japonicus Green, 1439
Chilocorus kuwanae Silvestri, 1439
methyl jasmonate, 1439
volatiles, 1439
plant morphology,
density dependent, 962
environmental conditions, 962
fertilization, 962
reproduction, 962
plant pathogens,
Cephus cinctus, 387
insect pathogens, 387
Triticum aestivum, 387
wheat, 387
plant quality,
elevational gradient, 1161
herbivory, 1161
phenolics, 1161
plant–herbivore interaction,
Aucuba japonica, 404
gall insect, 404
physical defense, 404
plant–herbivore interaction, 404
susceptible period, 404
plant–insect dynamics,
herbivore guild, 667
physiological structure, 667
plant–insect dynamics, 667
productivity, 667
resource competition, 667
plant–insect interactions,
angiosperm volatiles, 1189
Engelmann spruce weevil, 1189
nonhost volatiles, 1189
plant–insect interactions, 1189
spatial point processes, 1189
plant–insect–virus interactions,
Barley yellow dwarf virus epidemiology, 836
bird cherry-oat aphid, 836
plant–insect–virus interactions, 836
transgenic virus resistance, 836
virus-induced volatiles, 836
plasticity,
learning, 803
memory, 803
social wasps, 803
Plecoptera,
Aeromonas salmonicida, 1022
disease-pathogen dynamics, 1022
insect feeding assay, 1022
trophic interactions, 1022
Plutella xylostella,
cold tolerance, 524
overwintering, 524
pest management, 524
Polistinae,
Caatinga, 1613
community ecology, 1613
species richness, 1613
pollination syndromes,
antagonism, 741
Apocynaceae, 741
mutualism, 741
nectar, 741
pollinator performance,
Hoplitis, 1293
Megachile, 1293
Osmia, 1293
resource limitation, 1293
pollinator visitation,
environmental factors, 1480
Medicago sativa L., 1480
wild bee, 1480
polymerase chain reaction,
Acalymma vittatum, 395
Erwinia tracheiphila, 395
frass, 395
vector–pathogen interactions, 395
polymerase chain reaction-restriction fragment length polymorphism,
Austrocedrus chilensis, 505
Cinara cupressi, 505
Cinara tujafilina, 505
cytochrome oxidase I, 505
ponderosa pine,
competition, 639
Dendroctonus brevicomis, 639
Dendroctonus frontalis, 639
population dynamics, 639
poppy stem gall wasp,
biological control, 723
endophyte, 723
systemic protection, 723
Timaspis papaveris, 723
population abundance,
Frankliniella australis, 333
Frankliniella occidentalis, 333
Orius, 333
predation, 333
population dispersion,
invasive species, 1400
larval behavior, 1400
Olea europea, 1400
population dynamics, 639, 1205, 1716
Acrosternum hilare, 1716
black light trap, 1716
competition, 639
degree-day, 1716
Dendroctonus brevicomis, 639
Dendroctonus frontalis, 639
flight activity, 1716
Frankliniella fusca, 1205
Frankliniella occidentalis, 1205
Frankliniella tritici, 1205
ponderosa pine, 639
within-plant aggregation, 1205

population genetic structure,
Aphis glycines, 1301
microsatellites, 1301
soybean aphid, 1301
population genetics,
dispersal, 1312
gene flow, 1312
microsatellites, 1312
Ostrinia nubilalis, 1312
population model,
colonization process, 1575
onion, 1575
Thrips tabaci, 1575
positive interaction,
Anisopteromalus calandrae, 121
Aspergillus candidus, 121
Aspergillus niger, 121
Sitophilus oryzae, 121
postrelease evaluation,
dispersal, 1145
herbivory, 1145
invasive species, 1145
weed biological control, 1145
potato,
aphid behavior, 1429
emigration, 1429
Myzus persicae, 1429
virus-induced volatiles, 1429
powderpost beetle,
developmental period, 478
life cycle, 478
thermal constants, 478
thermal requirements, 478
precipitation, 307, 1618
ant–aphid interaction, 1618
damage, 307
foraging, 1618
phenology, 307
temperature, 307
voltinism, 307
precision pest management,
Fragaria spp., 137
spider mites, 137
visible and near infrared spectra, 137
predation,
Frankliniella australis, 333
Frankliniella occidentalis, 333
Orius, 333
population abundance, 333
predator,
Anura, 856
diving beetle, 856
fish, 856
Odonata, 856
predator–prey interaction,
biological control, 715
feeding behavior, 715
phytophagous mites, 715
predator–prey interaction, 715
predatory potential, 715
predatory potential,
biological control, 715
feeding behavior, 715
phytophagous mites, 715
predator–prey interaction, 715
prediction,
Choristoneura conflictana, 459
Malacosoma disstria, 459
monitor, 459
sex pheromone traps, 459
preoviposition behaviors,
Agral 90, 1387
antixenosis, 1387
pea leafminer, 1387
Sylgard 309, 1387
prevalence,
bacterial symbionts, 1069
distribution, 1069
Frankliniella occidentalis, 1069
prey quality,
Aphidius colemani, 153
aphids, 153
Coccinella undecimpunctata, 153
Myzus persicae, 153
primary attraction,
Cerambycidae, 797
host selection, 797
Scolytinae, 797
semiochemicals, 797
productivity,
herbivore guild, 667
physiological structure, 667
plant–insect dynamics, 667
resource competition, 667
protease inhibitor,
bacteria symbionts, 1078
pest control, 1078
proteases, 1078
16S rRNA, 1078
proteases,
bacteria symbionts, 1078
pest control, 1078
protease inhibitor, 1078
16S rRNA, 1078
Pseudonocardia,
Attini, 78
Escovopsis, 78
Leucoagaricus, 78
symbiosis, 78
psocids,
endosymbiont, 1365
life table parameters, 1365
thermal treatment, 1365
Pterostichus melanarius,
Cylindroiulus caeruleocinctus, 1106
integrated pest management, 1106
pupal predation,
autumnal moth, 627
sub-arctic birch forest, 627
winter moth, 627

R

(*R*)-3-hydroxyhexan-2-one,
longhorned beetle, 1462
mating behavior, 1462
Mojave desert, 1462
sex pheromone, 1462
ragweed beetle,
age, 235
dispersal potential, 235
flight mill, 235
photoperiod, 235
rangeland,
degree-day, 1521
grasshopper development, 1521
phenology, 1521
raspberry weevil,
Aegorhinus superciliosus, 192
contact pheromone, 192
mating behavior, 192
raster map,
brown stink bug, 1038
distribution, 1038
southern green stink bug, 1038
realized heritability,
beetle, 266
bidirectional selection, 266
exotic insect, 266
repeatability, 266
red turpentine beetle,
bark beetle phermone, 472
cis-verbenol, 472
frontalin, 472
trans-verbenol, 472
relative abundance,
biological control, 539
phenology, 539
sampling, 539
Solenopsis invicta, 539
remote sensing,
landscape ecology, 551
Solenopsis invicta, 551
repeatability,
beetle, 266
bidirectional selection, 266
exotic insect, 266
realized heritability, 266
reproduction, 962, 1061, 1283, 1496
Bactrocera oleae, 1496
C. transversoguttata, 1283
Coccinella septempunctata, 1283
density dependent, 962
development, 1061
egg maturation, 1496
egg size, 1283
egg viability, 1283
environmental conditions, 962
fertilization, 962
high temperature, 1496
intrinsic rate of increase, 1061
meta-analysis, 1061
plant morphology, 962
Russian wheat aphid, 1061
survival, 1496
reproductive biology,
baiting, 35
castes, 35
larvae, 35

subterranean termites, 35
reproductive capacity,
diets, 561
dispersal, 561
phenotypic plasticity, 561
small hive beetle, 561
reproductive maturity,
diapause, 569
locust, 569
Madagascar, 569
wing color, 569
resistance frequency,
Bt cotton, 1336
Cry1Ac, 1336
Helicoverpa armigera, 1336
resource competition,
herbivore guild, 667
physiological structure, 667
plant–insect dynamics, 667
productivity, 667
resource limitation,
Hoplitis, 1293
Megachile, 1293
Osmia, 1293
pollinator performance, 1293
restoration,
conservation management, 325
ecological disturbance, 325
mediterranean ecosystems, 325
review,
biosafety, 293
laboratory experiments, 293
natural enemies, 293
transgenic plants, 293
Rhagoletis pomonella,
Braconidae, 588
co-evolution, 588
North America, 588
Tephritidae, 588
Rhopalosiphum padi,
competition, 26
elevated CO_2, 26
Schizaphis graminum, 26
Sitobion avenae, 26
rice,
endured resistance, 1174
fall armyworm, 1174
jasmonic acid, 1174
Oryza sativa, 1174
Spodoptera frugiperda, 1174
rice stem borer,
Chilo suppressalis, 260
cold hardiness, 260
diapause, 260
glycerol, 260
risk analysis,
biological control, 2
insect demography, 2
multiple decrement life table, 2
natural enemies, 2
risk assessment,
Brazilian peppertree, 1642
Christmas berry, 1642
natural enemies, 1642
risk of parasitism,
agricultural landscapes, 380
biological control, 380
blueberry, 380
cottony cushion scale, 380
16S rRNA,
bacteria symbionts, 1078
pest control, 1078
protease inhibitor, 1078
proteases, 1078
Russian wheat aphid,
development, 1061
intrinsic rate of increase, 1061
meta-analysis, 1061
reproduction, 1061
ruthenium red dye,
integrated pest management, 1168
southern green stink bug, 1168

S

sacred groves,
edge effects, 43
fragmentation, 43
nonmetric multidimensional scaling, 43
West Africa, 43
sampling, 539, 1555
automated trapping, 1555
biological control, 539
flight behavior, 1555
Lymantria dispar, 1555
phenology, 539, 1555
relative abundance, 539
Solenopsis invicta, 539
sampling methods,
Formicidae, 1724
inventory, 1724
Ohio, 1724
species richness, 1724
sampling plan design,
Diabrotica undecimpunctata undecimpuncta, 809
integrated pest management, 809
scouting programs, 809
strip spraying, 809
scale-dependent effects,
density dependence, 790
host-parasitoid, 790
Identificaticoffee agroecosystem, 790
olfactory cue, 790
Scelionidae,
parasitoid, 375
Pentatomidae, 375
Telenomus podisi, 375
Trissolcus basalis, 375
Schizaphis graminum,
competition, 26
elevated CO_2, 26
Rhopalosiphum padi, 26
Sitobion avenae, 26
Scolytinae,
Cerambycidae, 797
host selection, 797
primary attraction, 797
semiochemicals, 797
scouting programs,
Diabrotica undecimpunctata undecimpuncta, 809
integrated pest management, 809
sampling plan design, 809
strip spraying, 809
seasonality,
ant community, 317
citrus orchard, 317
dominance, 317
Mediterranean region, 317
secondary salinization,
arbovirus, 1585
community, 1585
dryland salinity, 1585
temporary water, 1585
semiochemicals, 797, 1259
Caribbean fruit fly, 1259
Cerambycidae, 797
1,4-diaminobutane, 1259
host selection, 797
olfaction, 1259
primary attraction, 797
Scolytinae, 797
synthetic attractants, 1259
Sesiidae,
apple, 1219
behavior, 1219
dogwood borer, 1219
Synanthedon scitula, 1219
seven-spotted lady beetle,
biological control, 708
functional response, 708
multicolored Asian lady beetle, 708
soybean aphid, 708
sex attraction,
Cacopsylla pyricola, 815
diel periodicity, 815
pear psylla, 815
sex determination,
fitness, 499
isogenic strain, 499
M factor, 499
sex pheromone,
longhorned beetle, 1462
mating behavior, 1462
Mojave desert, 1462
(*R*)-3-hydroxyhexan-2-one, 1462
sex pheromone traps,
Choristoneura conflictana, 459
Malacosoma disstria, 459
monitor, 459
prediction, 459
sex-biased parasitism,
coelopidae, 1608

silkworm,
food chain, 1276
heavy metals, 1276
lead, 1276
mulberry, 1276
Sitobion avenae,
competition, 26
elevated CO_2, 26
Rhopalosiphum padi, 26
Schizaphis graminum, 26
Sitophilus oryzae,
Anisopteromalus calandrae, 121
Aspergillus candidus, 121
Aspergillus niger, 121
positive interaction, 121
small hive beetle,
diets, 561
dispersal, 561
phenotypic plasticity, 561
reproductive capacity, 561
snow cover,
Aphthona flea beetles, 1539
chill degree-days, 1539
low temperature, 1539
overwintering, 1539
social form,
Argentina, 198
biological control, 198
fire ants, 198
parasitoid, 198
social wasps,
learning, 803
memory, 803
plasticity, 803
Solanaceae,
development, 1661
fecundity, 1661
oviposition, 1661
survival, 1661
Solanum americanum,
Anthonomus eugenii, 216
Capsicum annuum, 216
host plant volatiles, 216
Solanum melongena, 216
Solanum melongena,
Anthonomus eugenii, 216
Capsicum annuum, 216
host plant volatiles, 216
Solanum americanum, 216
solar UV radiation,
habitat microenvironment, 920
Neoseiulus, 920
Tetranychus, 920
Solenopsis geminata,
abdomen length, 879
alkaloids, 879
body length, 879
venom volume, 879
Solenopsis invicta, 539, 551, 1241
biological control, 539
Coptotermes formosanus, 1241
foraging, 1241
integrated pest management, 1241
landscape ecology, 551
nesting, 1241
phenology, 539
relative abundance, 539
remote sensing, 551
sampling, 539
solitary wasps,
Arribes del Duero Natural Park, 608
biogeography, 608
diversity, 608
Mediterranean region, 608
southern green stink bug, 103, 1038, 1168
brown stink bug, 1038
Buchnera, 103
distribution, 1038
Heteroptera, 103
integrated pest management, 1168
raster map, 1038
ruthenium red dye, 1168
symbiosis, 103
soybean aphid, 110, 708, 1301
Aphis glycines, 1301
Arsenophonus, 110
bacterial symbiont, 110
biological control, 708
Buchnera aphidicola, 110
functional response, 708
microsatellites, 1301
multicolored Asian lady beetle, 708
population genetic structure, 1301
seven-spotted lady beetle, 708
spatial analysis of distance indices,
foraging behavior, 183
marginal value theorem, 183
spatial point processes,
angiosperm volatiles, 1189
Engelmann spruce weevil, 1189
nonhost volatiles, 1189
plant–insect interactions, 1189
spatial scales,
biological control, 365
host–parasitoid interaction, 365
Lycopersicon esculentum, 365
species diversity,
Argiope trifasciata, 1595
Cyclosa turbinata, 1595
Mangora gibberosa, 1595
Neoscona arabesca, 1595
species richness, 1613, 1724
Caatinga, 1613
community ecology, 1613
Formicidae, 1724
inventory, 1724
Ohio, 1724
Polistinae, 1613
sampling methods, 1724
spider mites,
Fragaria spp., 137
precision pest management, 137
visible and near infrared spectra, 137
spiders,
carabid beetles, 944
corn, 944
genetically modified organisms, 944
Orius, 944
spillover,
agroecology, 1028
agroecosystem, 1028
cross-boundary, 1028
Great Basin, 1028
Spodoptera frugiperda, 1086, 1174, 1546
context-dependent interactions, 1086
cytochrome oxidase, 1546
endured resistance, 1174
Epichloë, 1086
fall armyworm, 1174
food choice, 1086
haplotype, 1546
jasmonic acid, 1174
Neotyphodium, 1086
Noctuidae, 1546
Oryza sativa, 1174
rice, 1174
squash,
Bemisia argentifolii, 442
intercropping, 442
living mulch, 442
phytotoxemia, 442
stable fly,
house fly, 700
Muscidifurax raptorellus, 700
Tachinaephagus zealandicus, 700
Trichopria nigra, 700
starvation,
allocation, 1379
foraging, 1379
Helicoverpa zea, 1379
nutrition, 1379
Steinernema,
Beauveria bassiana, 1126
Heterorhabdtis, 1126
inundative biological control, 1126
Metarhizium anisopliae, 1126
stress,
biological invasion, 484
nest site selection, 484
parasitoid, 484
temperature, 484
Striacosta albicosta,
biological invasion, 274
NADH dehydrogenase 1, 274
Zea mays, 274

strip spraying,
Diabrotica undecimpunctata undecimpuncta, 809
integrated pest management, 809
sampling plan design, 809
scouting programs, 809
sub-arctic birch forest,
autumnal moth, 627
pupal predation, 627
winter moth, 627
subalpine,
bumble bee, 748
flower density, 748
food availability, 748
foraging distance, 748
subterranean termites,
baiting, 35
castes, 35
larvae, 35
reproductive biology, 35
sugar content,
contaminant pest, 1563
Harmonia axyridis, 1563
modeling, 1563
trap catching, 1563
supercooling capacity,
Chilo suppressalis, 1182
overwintering, 1182
paddy rice, 1182
water-oat, 1182
surface-floating pupal exuviae,
in situ enclosures, 250
instantaneous growth rate, 250
winter active, 250
survival, 1485, 1496, 1661
Bactrocera oleae, 1496
Cicadellidae, 1485
development, 1661
egg maturation, 1496
fecundity, 1661
feeding, 1485
glassy-winged sharpshooter, 1485
high temperature, 1496
oviposition, 1661
reproduction, 1496
Solanaceae, 1661
temperature, 1485
vector, 1485
susceptible period,
Aucuba japonica, 404
gall insect, 404
physical defense, 404
plant–herbivore interaction, 404
swarming,
chronological reproductive isolation, 1697
flight phenology, 1697
Macrotermes, 1697
sweet cherry,
egg, 955
ovary, 955
tart cherry, 955
Sylgard 309,
Agral 90, 1387
antixenosis, 1387
pea leafminer, 1387
preoviposition behaviors, 1387
symbiosis, 67, 78, 103, 1096
artificial medium, 1096
Attini, 78
bark beetles, 67
biological invasion, 1096
Buchnera, 103
context dependency, 67
cooperative breeding, 1096
Escovopsis, 78
Heteroptera, 103
insect agriculture, 1096
insects, 67
Leucoagaricus, 78
microorganisms, 67
Pseudonocardia, 78
southern green stink bug, 103
Synanthedon scitula, 530, 1219
apple, 530, 1219
behavior, 1219
dogwood borer, 1219
forest, 530
ornamentals, 530
pheromone trapping, 530
Sesiidae, 1219
synchronous display,
climate change, 1505
degree-day, 1505
firefly, 1505
phenology, 1505
synthetic attractants,
Caribbean fruit fly, 1259
1,4-diaminobutane, 1259
olfaction, 1259
semiochemicals, 1259
systemic protection,
biological control, 723
endophyte, 723
poppy stem gall wasp, 723
Timaspis papaveris, 723

T

Tachinaephagus zealandicus,
house fly, 700
Muscidifurax raptorellus, 700
stable fly, 700
Trichopria nigra, 700
tall fescue,
ergot alkaloids, 996
Neotyphodium, 996
trophic interactions, 996
tamarisk biocontrol,
Diorhabda elongata, 1373
host choice, 1373
introduced species, 1373
larval performance, 1373
Tarsonemus,
fungal transmission, 143
ophiostomatoid fungi, 143
phoresy, 143
vector, 143
tart cherry,
egg, 955
ovary, 955
sweet cherry, 955
Telenomus isis,
East African stemborer species, 904
host acceptability and suitability, 904
life table, 904
temperature and humidity requirements, 904
Telenomus podisi,
parasitoid, 375
Pentatomidae, 375
Scelionidae, 375
Trissolcus basalis, 375
temperature, 18, 307, 484, 633, 930, 1485
age-dependent developmental rate, 18
ammonia, 633
biological invasion, 484
biosolid cake, 633
black soldier fly, 930
Cicadellidae, 1485
damage, 307
development, 930
egg phenology, 18
feeding, 1485
fly production, 633
glassy-winged sharpshooter, 1485
Hermetia illucens, 930
nest site selection, 484
parasitoid, 484
phenology, 18, 307
precipitation, 307
stress, 484
survival, 1485
vector, 1485
voltinism, 307
temperature and humidity requirements,
East African stemborer species, 904
host acceptability and suitability, 904
life table, 904
Telenomus isis, 904
temporary water,
arbovirus, 1585
community, 1585
dryland salinity, 1585
secondary salinization, 1585
Tephritidae, 116, 588
Braconidae, 588
co-evolution, 588
ecology, 116
host fruit, 116
microorganisms, 116
North America, 588
Rhagoletis pomonella, 588
trapping, 116

Tetranychus,
habitat microenvironment, 920
Neoseiulus, 920
solar UV radiation, 920
Therioaphis trifolii,
Acyrthosiphon pisum, 1347
Aphis glycines, 1347
Aphis gossypii, 1347
Phaseolus vulgaris, 1347
thermal constants,
developmental period, 478
life cycle, 478
powderpost beetle, 478
thermal requirements, 478
thermal requirements,
developmental period, 478
life cycle, 478
powderpost beetle, 478
thermal constants, 478
thermal threshold,
codling moth, 885
Cydia pomonella, 885
development, 885
linear and nonlinear models, 885
thermal treatment,
endosymbiont, 1365
life table parameters, 1365
psocids, 1365
Thrips tabaci,
colonization process, 1575
onion, 1575
population model, 1575
Timaspis papaveris,
biological control, 723
endophyte, 723
poppy stem gall wasp, 723
systemic protection, 723
Torymus,
community interactions, 417
endophyte, 417
gypsy moth, 417
lesser chestnut weevil, 417
trans-verbenol,
bark beetle pheromone, 472
cis-verbenol, 472
frontalin, 472
red turpentine beetle, 472
transect,
canonical correlation analysis, 1600
canopy trap, 1600
distributional pattern, 1600
elevational distribution, 1600
transgenic plants,
biosafety, 293
laboratory experiments, 293
natural enemies, 293
review, 293
transgenic virus resistance,
Barley yellow dwarf virus epidemiology, 836
bird cherry-oat aphid, 836
plant–insect–virus interactions, 836
virus-induced volatiles, 836
trap catching,
contaminant pest, 1563
Harmonia axyridis, 1563
modeling, 1563
sugar content, 1563
trap crop,
Acrosternum hilare, 450
Euschistus servus, 450
Glycine max, 450
Nezara viridula, 450
trap design,
Anoplophora glabripennis, 1745
Cerambycidae, 1745
male-produced pheromone, 1745
mate-finding, 1745
trap tree, 1226, 1668
Agrilus planipennis, 1226
detection methods, 1668
emerald ash borer, 1226
Fraxinus, 1226
girdled trees, 1668
Manuka oil, 1668
methyl jasmonate, 1668
tree stress, 1226
Trapa natans,
host preference and performance, 1653
insect–plant interaction, 1653
invasion, 1653
Nuphar lutea, 1653
trapping,
ecology, 116
host fruit, 116
microorganisms, 116
Tephritidae, 116
tree disease,
insect/fungus associations, 356
ophiostomatoid fungi, 356
tree mortality,
bark beetles, 1197
Dendroctonus rhizophagus, 1197
host preference, 1197
pine seedlings, 1197
tree stress,
Agrilus planipennis, 1226
emerald ash borer, 1226
Fraxinus, 1226
trap tree, 1226
Trichogramma cacoeciae,
host age, 1513
host density, 1513
Lobesia botrana, 1513
UV treatment, 1513
Trichogramma,
biological control, 677
dispersal, 677
European corn borer, 677
integrated pest management, 677
Trichopria nigra,
house fly, 700
Muscidifurax raptorellus, 700
stable fly, 700
Tachinaephagus zealandicus, 700
Trichosirocalus horridus,
biological control of weeds, 731
Cirsium, 731
ecological risk assessment, 731
weed control, 731
Trissolcus basalis,
parasitoid, 375
Pentatomidae, 375
Scelionidae, 375
Telenomus podisi, 375
Triticum aestivum,
Cephus cinctus, 387
insect pathogens, 387
plant pathogens, 387
wheat, 387
trophic interactions, 996, 1022
Aeromonas salmonicida, 1022
disease-pathogen dynamics, 1022
ergot alkaloids, 996
insect feeding assay, 1022
Neotyphodium, 996
Plecoptera, 1022
tall fescue, 996

U

ungulates,
arachnids, 985
breeds, 985
grazing pressure, 985
insects, 985
urban ecology,
ant nest, 600
nest-site limitation, 600
UV treatment,
host age, 1513
host density, 1513
Lobesia botrana, 1513
Trichogramma cacoeciae, 1513

V

vector, 143, 1485
Cicadellidae, 1485
feeding, 1485
fungal transmission, 143
glassy-winged sharpshooter, 1485
ophiostomatoid fungi, 143
phoresy, 143
survival, 1485
Tarsonemus, 143
temperature, 1485
vector biology,
disease management, 345
epidemiology, 345
vector control,
Aedes vigilax, 1013
Culex annulirostris, 1013
density dependence, 1013
larval habitats, 1013

vector–pathogen interactions,
Acalymma vittatum, 395
Erwinia tracheiphila, 395
frass, 395
polymerase chain reaction, 395
vector–pathogen interactions, 395
venom volume,
abdomen length, 879
alkaloids, 879
body length, 879
Solenopsis geminata, 879
virus-induced volatiles, 836, 1429
aphid behavior, 1429
Barley yellow dwarf virus epidemiology, 836
bird cherry-oat aphid, 836
emigration, 1429
Myzus persicae, 1429
plant–insect–virus interactions, 836
potato, 1429
transgenic virus resistance, 836
visible and near infrared spectra,
Fragaria spp., 137
precision pest management, 137
spider mites, 137
visual cues,
natural enemies, 159
olfactometer, 159
phenylacetaldehyde, 159
western tarnished plant bug, 159
volatile compounds,
crucifer flea beetle, 1467
olfactometer bioassay, 1467
volatiles,
Ceroplastes japonicus Green, 1439
Chilocorus kuwanae Silvestri, 1439
methyl jasmonate, 1439
α-pinene, 1439
voltinism,
damage, 307
phenology, 307
precipitation, 307
temperature, 307

W

water sources,
Diptera, 657
forest management, 657
intensive forestry, 657
pine plantation, 657
water-oat,
Chilo suppressalis, 1182
overwintering, 1182
paddy rice, 1182
supercooling capacity, 1182
weed biological control,
dispersal, 1145
herbivory, 1145
invasive species, 1145
postrelease evaluation, 1145
weed control,
biological control of weeds, 731
Cirsium, 731
ecological risk assessment, 731
Trichosirocalus horridus, 731
weevil, 781, 870, 1423
agave, 1423
attractant, 870
electroantennogram, 870
ethologic control, 781
gas chromatographic-electroantennographic detection, 870
integrated pest management, 781
2-methyl-4-octanone, 1423
olfactometric behavior, 781
pheromone, 870
pheromone release, 1423
West Africa,
edge effects, 43
fragmentation, 43
nonmetric multidimensional scaling, 43
sacred groves, 43
western corn rootworm,
Bt corn, 176
dispersal, 176
insect resistance management, 176
western tarnished plant bug,
natural enemies, 159
olfactometer, 159
phenylacetaldehyde, 159
visual cues, 159
wheat,
Cephus cinctus, 387
insect pathogens, 387
plant pathogens, 387
Triticum aestivum, 387
white-footed mouse,
blacklegged tick, 977
Japanese barberry, 977
Lyme disease, 977
white-tailed deer, 977
white-tailed deer,
blacklegged tick, 977
Japanese barberry, 977
Lyme disease, 977
white-footed mouse, 977
wild bee,
environmental factors, 1480
Medicago sativa L., 1480
pollinator visitation, 1480
wing color,
diapause, 569
locust, 569
Madagascar, 569
reproductive maturity, 569
winter active,
in situ enclosures, 250
instantaneous growth rate, 250
surface-floating pupal exuviae, 250
winter moth,
autumnal moth, 627
pupal predation, 627
sub-arctic birch forest, 627
winter squash,
Apis mellifera, 207
Bombus spp., 207
multispecies interactions, 207
Peponapis pruinosa, 207
within-plant aggregation,
Frankliniella fusca, 1205
Frankliniella occidentalis, 1205
Frankliniella tritici, 1205
population dynamics, 1205
within-plant patterns,
Aphis craccivora, 1765
cowpea aphid, 1765
developmental threshold, 1765
Wolbachia,
cytoplasmic incompatibility, 1626
Laodelphax striatellus, 1626
microinjection, 1626
Nilaparvata lugens, 1626

Y

Y-tube bioassay,
electroantennogram, 861
gas chromatography-mass spectrometry, 861
headspace analysis, 861
highbush blueberry, 861
Yucca filamentosa,
Florida, 1211
host plant apparency, 1211
larval performance, 1211
oviposition preference, 1211

Z

Zea mays,
biological invasion, 274
NADH dehydrogenase 1, 274
Striacosta albicosta, 274